




COLONOSCOPY:

WHAT TO EXPECT

BEFORE, DURING, AND AFTER



CONGRATULATIONS!


You've made a smart decision to help take care of your health by undergoing a colonoscopy. A colonoscopy is a routine, safe, and relatively painless procedure that will allow your doctor to see the lining of your **colon**.

More importantly, a screening colonoscopy is the best way to detect and prevent colorectal cancer (CRC), the second leading cause of death from cancer in the United States. A colonoscopy can also help doctors diagnose certain other gastrointestinal disorders.

This brochure was designed to help you better understand the colonoscopy procedure: what you'll need to do to prepare, what you can expect during the procedure, and how you can help ensure a successful exam. It describes the entire process and provides useful hints to help make the colonoscopy process more comfortable and easier for you.

To help make this information clear, we've provided a glossary with definitions of certain words on pages 22–23. Any **bold** word on the following pages is defined in the glossary.


Anatomy of the lower digestive tract



UNDERSTANDING BOWEL PREP

Before you go to your doctor's office for your **colonoscopy**, you'll need to take a bowel prep. A bowel prep is a prescription medication that will prepare your **colon** for viewing by removing solid waste from your digestive tract. In other words, it causes episodes of diarrhea occurring over several hours, so that the **colon** is empty at the time of your **colonoscopy** and the doctor can see the lining of the **colon** clearly.

The complete bowel prep regimen includes a special diet as well as a prescription medication. You'll need to start the bowel prep process a full day before going in for your procedure, or as directed by your healthcare provider.



BOWEL PREP OPTIONS

Together, you and your healthcare provider will choose the prep that is right for you. Make sure your healthcare provider knows in advance about any medications you are taking, as well as any past or ongoing medical conditions, including heart, kidney, liver, and nervous system problems.

Preps come in several forms:



Liquid preps require you to drink a prep solution and vary in volume and taste and are often accompanied by clear liquids for **hydration**



Tablet preps are taken as a series of pills, accompanied by clear liquids for **hydration**



Once your prep is selected, it is important to follow your healthcare provider's instructions exactly and completely. How successfully your bowel is prepped determines how well your doctor can see the **colon** lining and help detect potential problems.

PREPPING WITH PREPOPIK


Prepopik is a prescription medicine used by adults to clean the **colon** before a **colonoscopy**. Prepopik cleanses your **colon** by causing you to have diarrhea. Cleansing your **colon** helps your doctor see the inside of your **colon** more clearly during your **colonoscopy**.

Prepopik is the bowel prep with the lowest volume of prep solution, which is an orange-flavored powder dissolved in water. Prepopik also allows you to drink any approved clear liquids at your own pace, to maintain **hydration** during the prep process.

Do not take Prepopik if your healthcare provider has told you that you have:

- Serious kidney problems
- A blockage in your intestine (bowel obstruction)
- An opening in the wall of your stomach or intestines (bowel perforation)
- A very dilated intestine (toxic megacolon)
- Problems with the emptying of food and fluid from your stomach (gastric retention)
- An allergy to any of the ingredients in Prepopik

Please see pages 20–21 for additional Important Safety Information and enclosed Medication Guide.

 **Prepopik**[™]
(sodium picosulfate, magnesium oxide, and
anhydrous citric acid) for oral solution
10 mg/3.5 g/12 g per packet



How is Prepopik different from other therapies?

Prepopik allows you to drink less prep solution and stay hydrated with the clear liquids of your choice.

How is Prepopik taken? Your Prepopik dose is taken in 2 parts, each requiring only 5 ounces of prep solution, followed by additional clear liquids of your choice.

How does Prepopik work?

Prepopik causes diarrhea that, over time, cleans the solid waste from your **colon**.

How quickly does Prepopik work?

Each patient reacts to Prepopik differently. The effects of Prepopik may be felt right away or they may take some time. It is important for you to have regular access to a bathroom as soon as you begin taking Prepopik.

Where can I find more instructions for taking Prepopik?

The full instructions for prepping with Prepopik are available in the official Medication Guide.



Ask your healthcare provider if Prepopik is right for you.

EATING AND DRINKING DURING YOUR PREP

Starting the day before your procedure, you will only be able to eat clear foods (gelatin, broth, etc.) and liquids. Be sure to get exact instructions from your healthcare provider about when you need to stop eating solids, what you can and cannot drink, and how often you should drink during the bowel prep.

Stay hydrated

A major part of prep success is staying hydrated. The prep process removes a lot of water from your body, and it needs to be replaced.

Some preps include clear fluids as part of the regimen to help with proper **hydration**. Regardless of what prep you use, be sure to follow your healthcare provider's instructions for **hydration** carefully and completely.

Clear liquids

Always confirm with your healthcare provider before making any other choices.

Examples of clear liquids your healthcare provider may recommend:

- Water
- Clear broth
- Apple juice
- White cranberry juice
- White grape juice
- Ginger ale
- Plain gelatin
- Frozen juice bars



DO NOT drink any of the following:

- **NO** red or purple liquids
- **NO** liquids that you can't see through
- **NO** alcoholic beverages



Drinking approved clear liquids during prep is very important to keep you hydrated and to help the prep work properly.



WHAT TO EXPECT WHILE YOU PREP

Once you begin drinking the prep, it's hard to know when it will start working. Different people have different reactions. When the effects do kick in, they will be urgent and frequent. You'll need regular access to a bathroom from the time you start the prep, to the time you leave for your **colonoscopy** appointment. The effects of the prep may last for a few hours after you finish it.



Bowel cleansing is never a comfortable process. You should talk to your healthcare provider to understand what symptoms are normal side effects and what may signal an allergic reaction or dehydration that requires immediate attention.

Tell your healthcare provider right away if you have any of these symptoms of a loss of too much body fluid (dehydration):

- Vomiting
- Urinating less often than normal
- Dizziness
- Headache



TIPS FOR PREP SUCCESS

i

Inform yourself

Before starting, be sure that you clearly understand:

- All the steps of the bowel prep process
- All of your healthcare provider's instructions, particularly when to start and finish the prep
- When, and what, you can and cannot eat or drink before, during, and after the prep and **colonoscopy**



Stay comfortable

Once you start drinking the prep, you'll be spending a lot of time in the bathroom.

- Bring a book or magazine to read
- Bring a handheld game to play
- Use moistened wipes or water spray instead of toilet paper to avoid irritation
- Wear pants that are easy to take on and off repeatedly



Make sure you finish

It's important to finish the entire prep so your doctor can see your **colon** clearly. If you don't finish, your **colon** may not be cleansed completely. The **colonoscopy** may take longer than expected and, if the prep is really unsuccessful, the doctor may stop in the middle and ask you to reschedule. This will mean going through the prep process and procedure all over again.



WHAT TO EXPECT DURING YOUR COLONOSCOPY



Before the procedure

When you arrive, there will likely be paperwork to fill out. Afterward, you will change into a gown and be fitted with an **IV** and other monitoring devices that will keep track of your vital signs. The **IV** will allow the doctor and his or her staff to administer **sedation** throughout the procedure.



During the procedure

For the procedure itself, you'll lay on your side, and once the **sedation** takes effect, you won't remember much else.

While you are sedated, the doctor will insert a long, hose-like instrument called a **colonoscope** into your **anus** and through the natural cavity of your **rectum** and **colon**. The **colonoscope** is fitted with a light, a camera, and other instruments that the doctor will use to examine your **colon** for:

- **Inflammation** (swelling and/or redness)
- Other precancerous **lesions** and potential cancer
- **Polyps**
- Other signs of digestive problems

The procedure also allows the doctor to remove **polyps** before they become cancerous and take other small samples, or **biopsies**, for lab analysis. Remember that there are no nerve endings in the lining of the **colon**. **Biopsies** and **polyp** removal should be a relatively painless process. If you have any questions about your procedure, contact your healthcare provider.



After the procedure

After the **colonoscopy**, you will be taken to a recovery room where you can rest until the **sedation** partially wears off. Once you are awake, your healthcare provider will speak to you about the results of the **colonoscopy** and you will receive a written report with photos and initial results from the exam. Any **polyps** or other **lesions** that are removed or **biopsied** will be sent to a lab for analysis.

You may feel some bloating from the procedure and pass some gas. These symptoms should subside within a few hours.

Getting home and taking it easy

A few days before the procedure, make sure to arrange for a friend or family member to escort you home. The effects of sedation make it unsafe for you to drive or find your way home alone.

Most doctors' offices will not allow you to be sedated until you provide proof of an escort. Be sure to arrange one in advance.

Don't plan on driving or operating any machinery until after you've had a full night's sleep. Once you get home:

- Plan on resting for the remainder of the day
- Ease back into eating after your **colonoscopy**. Start with a small, light meal and work gradually back up to your normal diet



If you have questions or concerns about the procedure itself or about your results, call your healthcare provider for more details and explanation.



Important Safety Information

- Do not take Prepopik if your healthcare provider has told you that you have serious kidney problems, a blockage in your intestine (bowel obstruction), an opening in the wall of your stomach or intestines (bowel perforation), a very dilated intestine (toxic megacolon), problems with the emptying of food and fluid from your stomach (gastric retention), or an allergy to any of the ingredients in Prepopik
- Tell your healthcare provider right away if you have any of these symptoms of a loss of too much body fluid (dehydration) while taking Prepopik: vomiting that prevents you from keeping down the additional prescribed amounts of clear liquids that you must drink after taking Prepopik, dizziness, urinating less often than normal, or headache

- Prepopik and other bowel preparations can cause serious side effects, including serious loss of body fluid (dehydration) and changes in blood salts (electrolytes) in your blood. These changes can cause abnormal heartbeats that can cause death, seizures (this can happen even if you have never had a seizure), or kidney problems. Your chance of having fluid loss and changes in blood salts with Prepopik is higher if you have heart problems, have kidney problems, or take water pills or non-steroidal anti-inflammatory drugs (NSAIDs). The most common side effects of Prepopik include nausea, headache, and vomiting

GLOSSARY

Anus: The final opening at the end of the digestive tract

Biopsy: A tissue or fluid sample removed from the body for lab analysis

Colon: The part of the large intestine between the cecum and the **rectum**

Colonoscope: A long, hose-like instrument fitted with a light, a camera, and other attachments that doctors use to examine the **colon**

Colonoscopy: A routine, safe, and relatively painless procedure that allows doctors to literally see the lining of the **colon**

Hydration: The introduction of fluid into the body, normally through drinking

Inflammation: Any bodily response that causes swelling, pain, redness, and/or loss of function

IV: Short for intravenous line, a device used to administer fluids directly through the bloodstream

Lesion: Any change in the appearance of an organ or body part that suggests injury

Polyp: Any small mass of abnormal tissue that sticks out from the wall of a body part or organ

Rectum: The chamber of the large intestine directly above the **anus**

Sedation: Relaxed, sometimes unconscious state achieved with tranquilizing medications

Colonoscopy Checklist

Have you:



Gone over the bowel prep instructions with your doctor or someone from your doctor's office?



Filled your bowel prep prescription and prepared to take the prep?



Arranged for a friend or family member to take you to and from your colonoscopy?

Call your doctor's office if you have other questions or concerns



© 2012 Ferring B.V.
PREPOPIK™ is a trademark of Ferring B.V.
PREP_PEB_001_0812

Please see enclosed full Product Information.
For more information, visit www.prepopik.com.

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use PREPOPIK safely and effectively. See full prescribing information for PREPOPIK.

PREPOPIK (sodium picosulfate, magnesium oxide, and anhydrous citric acid) for oral solution

Initial U.S. Approval: 2012

-----INDICATIONS AND USAGE-----

PREPOPIK is a combination of sodium picosulfate, a stimulant laxative, and magnesium oxide and anhydrous citric acid which form magnesium citrate, an osmotic laxative, indicated for cleansing of the colon as a preparation for colonoscopy in adults (1)

-----DOSAGE AND ADMINISTRATION-----

- PREPOPIK, supplied as a powder, must be reconstituted with cold water right before its use (2.1, 2.2).
- Two dosing regimens, each requires two separate dosing times (2.1)
- “Split Dose” method is preferred method (2.3)
 - First dose: during evening before the colonoscopy
 - Second dose: next day, during the morning prior to the colonoscopy
- “Day Before” method is alternative method if “Split Dose” is not appropriate (2.4)
 - First dose: during afternoon or early evening before the colonoscopy
 - Second dose: 6 hours later during evening before colonoscopy
- Additional clear liquids (no solid food or milk) must be consumed after every dose in both dosing regimens (2.3, 2.4).

-----DOSAGE FORMS AND STRENGTHS-----

For oral solution: each of 2 packets contains 16.1 g of powder: 10 mg sodium picosulfate, 3.5 g magnesium oxide, and 12 g anhydrous citric acid (3)

-----CONTRAINDICATIONS-----

- Patients with severely reduced renal function (creatinine clearance less than 30 mL/minute (4)
- Gastrointestinal (GI) obstruction or ileus (4)
- Bowel perforation (4)
- Toxic colitis or toxic megacolon (4)
- Gastric retention (4)

-----WARNINGS AND PRECAUTIONS-----

- *Risk of fluid and electrolyte abnormalities, arrhythmia, seizures, and renal impairment:* Encourage adequate hydration, assess concurrent medications, and consider laboratory assessments prior to and after use (5.1, 5.2, 5.3, 5.4)
- *Risks in patients with renal insufficiency or patients taking concomitant medications that affect renal function:* Use caution, ensure adequate hydration and consider testing (5.3)
- *Mucosal ulcerations:* Consider potential for mucosal ulcerations when interpreting colonoscopy findings in patients with known or suspected inflammatory bowel disease (5.5)
- *Suspected GI obstruction or perforation:* Rule out diagnosis before administration (4, 5.6)
- *Patients at risk for aspiration:* Observe during administration (5.7)
- *Not for direct ingestion:* Dissolve and take with additional water (5.8)

-----ADVERSE REACTIONS-----

Most common adverse reactions (>1%) are nausea, headache and vomiting (abdominal bloating, distension, pain/cramping, and watery diarrhea not requiring an intervention were not collected) (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact Ferring at (1-888-FERRING (1-888-337-7464) or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

-----DRUG INTERACTIONS-----

- Drugs that increase risks due to fluid and electrolyte change (7.1)
- *Oral medication taken within 1 hour of start of each dosing:* Might not be properly absorbed (7.2)
- *Antibiotics:* Prior or concomitant use of antibiotics may reduce efficacy of PREPOPIK (7.3)

-----USE IN SPECIFIC POPULATIONS-----

Pregnancy: PREPOPIK should be used during pregnancy only if clearly needed (8.1)

See 17 for PATIENT COUNSELING INFORMATION

Revised: July 2012

FULL PRESCRIBING INFORMATION: CONTENTS*

- 1 INDICATIONS AND USAGE**
- 2 DOSAGE AND ADMINISTRATION**
- 3 DOSAGE FORMS AND STRENGTHS**
- 4 CONTRAINDICATIONS**
- 5 WARNINGS AND PRECAUTIONS**
 - 5.1 Serious Fluid and Serum Chemistry Abnormalities
 - 5.2 Seizures
 - 5.3 Use in Patients with Renal Impairment
 - 5.4 Cardiac Arrhythmias
 - 5.5 Colonic Mucosal Ulceration, Ischemic Colitis and Ulcerative Colitis
 - 5.6 Use in Patients with Significant Gastrointestinal Disease
 - 5.7 Aspiration
 - 5.8 Not for direct ingestion-dissolve and take with additional fluids
- 6 ADVERSE REACTIONS**
 - 6.1 Clinical Trials Experience
 - 6.2 Postmarketing Experience
- 7 DRUG INTERACTIONS**
 - 7.1 Drugs That May Increase Risks of Fluid and Electrolyte Abnormalities
 - 7.2 Potential for Altered Drug Absorption
 - 7.3 Antibiotics

- 8 USE IN SPECIFIC POPULATIONS**
 - 8.1 Pregnancy
 - 8.3 Nursing Mothers
 - 8.4 Pediatric Use
 - 8.5 Geriatric Use
 - 8.6 Renal Insufficiency
- 10 OVERDOSAGE**
- 11 DESCRIPTION**
- 12 CLINICAL PHARMACOLOGY**
 - 12.1 Mechanism of Action
 - 12.2 Pharmacodynamics
 - 12.3 Pharmacokinetics
- 13 NONCLINICAL TOXICOLOGY**
 - 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility
- 14 CLINICAL STUDIES**
- 16 HOW SUPPLIED/STORAGE AND HANDLING**
- 17 PATIENT COUNSELING INFORMATION and Medication Guide**

*Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

PREPOPIK™ (sodium picosulfate, magnesium oxide and anhydrous citric acid) for oral solution is indicated for cleansing of the colon as a preparation for colonoscopy in adults.

2 DOSAGE AND ADMINISTRATION

2.1 Dosing Overview

PREPOPIK, supplied as a powder, must be reconstituted with cold water right before its use [see *Dosage and Administration (2.2)*]. There are two dosing regimens, each requires two separate dosing times:

- The preferred method is the “Split Dose” method and consists of two separate doses: the first dose during the evening before the colonoscopy and the second dose the next day, during the morning prior to the colonoscopy [see *Dosage and Administration (2.3)*]
- The alternative method is the “Day Before” method and consists of two separate doses: the first dose during the afternoon or early evening before the colonoscopy and the second dose 6 hours later during the evening before the colonoscopy [see *Dosage and Administration (2.4)*].

Additional fluids must be consumed after every dose in both dosing regimens [see *Dosage and Administration (2.3, 2.4)*]. Instruct patients to consume only clear liquids (no solid food or milk) on the day before the colonoscopy up until 2 hours before the time of the colonoscopy. Instruct patients that if they experience severe bloating, distention, or abdominal pain following the first dose, delay the second dose until their symptoms resolve.

2.2 Reconstitution of the PREPOPIK Powder

- (a) Reconstitute the PREPOPIK powder right before each administration. Do not prepare the solution in advance.
- (b) Fill the supplied dosing cup with cold water up to the lower (5-ounce) line on the cup and pour in the contents of one packet of PREPOPIK powder.
- (c) Stir for 2 to 3 minutes. The reconstituted PREPOPIK solution may become slightly warm as the powder dissolves.

2.3 Split-Dose Dosing Regimen (Preferred Method)

The Split-Dose regimen is the preferred dosing method. Instruct patients to take two separate doses in conjunction with fluids, as follows:

- Take the first dose during the evening before the colonoscopy (e.g., 5:00 to 9:00 PM) followed by five 8-ounce drinks (upper line on the dosing cup) of clear liquids before bed. Consume clear liquids within 5 hours.
- Take second dose, the next day approximately 5 hours before the colonoscopy followed by at least three 8-ounce drinks of clear liquids before the colonoscopy. Consume clear liquids within 5 hours up until 2 hour before the time of the colonoscopy.

2.4 Day-Before Dosing Regimen (Alternative Method)

The Day-Before regimen is the alternative dosing method for patients for whom the Split-Dosing is inappropriate. Instruct patients to take two separate doses in conjunction with fluids, as follows:

- Take the first dose in the afternoon or early evening (e.g., 4:00 to 6:00 PM) before the colonoscopy followed by five 8-ounce drinks (upper line on the dosing cup) of clear liquids before the next dose. Consume clear liquids within 5 hours.
- Take the second dose approximately 6 hours later in the late evening (e.g., 10:00 PM to 12:00 AM), the night before the colonoscopy followed by three 8-ounce drinks of clear liquids before bed. Consume clear liquids within 5 hours.

3 DOSAGE FORMS AND STRENGTHS

For oral solution: each of the two packets contains 10 mg of sodium picosulfate, 3.5 grams of magnesium oxide, and 12.0 grams of anhydrous citric acid in 16.1 grams of powder

4 CONTRAINDICATIONS

PREPOPIK is contraindicated in the following conditions:

- Patients with severely reduced renal function (creatinine clearance less than 30 mL/minute) which may result in accumulation of magnesium [see *Warnings and Precautions* (5.3)]
- Gastrointestinal obstruction or ileus [see *Warnings and Precautions* (5.6)]
- Bowel perforation
- Toxic colitis or toxic megacolon
- Gastric retention
- An allergy to any of the ingredients in PREPOPIK

5 WARNINGS AND PRECAUTIONS

5.1 Serious Fluid and Serum Chemistry Abnormalities

Advise patients to hydrate adequately before, during, and after the use of PREPOPIK. Use caution in patients with congestive heart failure when replacing fluids. If a patient develops significant vomiting or signs of dehydration including signs of orthostatic hypotension after taking PREPOPIK, consider performing post-colonoscopy lab tests (electrolytes, creatinine, and BUN) and treat accordingly. Approximately 20% of patients in both arms (PREPOPIK, 2L of PEG + E plus two x 5-mg bisacodyl tablets) of clinical trials of PREPOPIK had orthostatic changes (changes in blood pressure and/or heart rate) on the day of colonoscopy. In clinical trials orthostatic changes were documented out to seven days post colonoscopy. [see *Adverse Reactions* (6.1, 6.2)]

Fluid and electrolyte disturbances can lead to serious adverse events including cardiac arrhythmias or seizures and renal impairment. Fluid and electrolyte abnormalities should be corrected before treatment with PREPOPIK. In addition, use caution when prescribing Prepopik for patients who have conditions or who are using medications that increase the risk for fluid and electrolyte disturbances or that may increase the risk of adverse events of seizure, arrhythmia, and renal impairment.

5.2 Seizures

There have been reports of generalized tonic-clonic seizures with the use of bowel preparation products in patients with no prior history of seizures. The seizure cases were associated with electrolyte abnormalities (e.g., hyponatremia, hypokalemia, hypocalcemia, and hypomagnesemia) and low serum osmolality. The neurologic abnormalities resolved with correction of fluid and electrolyte abnormalities.

Use caution when prescribing PREPOPIK for patients with a history of seizures and in patients at risk of seizure, such as patients taking medications that lower the seizure threshold (e.g., tricyclic antidepressants), patients withdrawing from alcohol or benzodiazepines, patients with known or suspected hyponatremia. [see *Adverse Reactions* (6.2)]

5.3 Use in Patients with Renal Impairment

As in other magnesium containing bowel preparations, use caution when prescribing PREPOPIK for patients with impaired renal function or patients taking concomitant medications that may affect renal function (such as diuretics, angiotensin converting enzyme inhibitors, angiotensin receptor blockers, or non-steroidal anti-inflammatory drugs). These patients may be at increased risk for renal injury. Advise these patients of the importance of adequate hydration before during and after the use of PREPOPIK. Consider performing baseline and post-colonoscopy laboratory tests (electrolytes, creatinine, and BUN) in these patients. In patients with severely reduced renal function (creatinine clearance < 30 mL/min), accumulation of magnesium in plasma may occur.

5.4 Cardiac Arrhythmias

There have been rare reports of serious arrhythmias associated with the use of ionic osmotic laxative products for bowel preparation. Use caution when prescribing PREPOPIK for patients at increased risk of arrhythmias (e.g., patients with a history of prolonged QT, uncontrolled arrhythmias, recent myocardial infarction, unstable angina, congestive heart failure, or cardiomyopathy). Pre-dose and post-colonoscopy ECGs should be considered in patients at increased risk of serious cardiac arrhythmias.

5.5 Colonic Mucosal Ulceration, Ischemic Colitis and Ulcerative Colitis

Osmotic laxatives may produce colonic mucosal aphthous ulcerations and there have been reports of more serious cases of ischemic colitis requiring hospitalization. Concurrent use of additional stimulant laxatives with PREPOPIK may increase this risk. The potential for mucosal ulcerations should be considered when interpreting colonoscopy findings in patients with known or suspected inflammatory bowel disease. [see *Adverse Reactions* (6.2)]

5.6 Use in Patients with Significant Gastrointestinal Disease

If gastrointestinal obstruction or perforation is suspected, perform appropriate diagnostic studies to rule out these conditions before administering PREPOPIK. Use with caution in patients with severe active ulcerative colitis.

5.7 Aspiration

Patients with impaired gag reflex and patients prone to regurgitation or aspiration should be observed during the administration of PREPOPIK. Use with caution in these patients.

5.8 Not for Direct Ingestion

Each packet must be dissolved in 5 ounces of cold water and administered at separate times according to the dosing regimen. Ingestion of additional water is important to patient tolerance. Direct ingestion of the undissolved powder may increase the risk of nausea, vomiting, dehydration, and electrolyte disturbances.

6 ADVERSE REACTIONS

6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in clinical trials of another drug and may not reflect the rates observed in practice.

In randomized, multicenter, controlled clinical trials, nausea, headache, and vomiting were the most common adverse reactions (>1%) following PREPOPIK administration. The patients were not blinded to the study drug. Since abdominal bloating, distension, pain/cramping, and watery diarrhea are known to occur in response to colon cleansing preparations, these effects were documented as adverse events in the clinical trials only if they required medical intervention (such as a change in study drug or led to study discontinuation, therapeutic or diagnostic procedures, met the criteria for a serious adverse event), or showed clinically significant worsening during the study that was not in the frame of the usual clinical course, as determined by the investigator.

PREPOPIK was compared for colon cleansing effectiveness with a preparation containing two liters (2L) of polyethylene glycol plus electrolytes solution (PEG + E) and two 5-mg bisacodyl tablets, all administered the day before the procedure. Table 1 displays the most common adverse reactions in Study 1 and Study 2 for the PREPOPIK Split-Dose and Day-Before dosing regimens, respectively, each as compared to the comparator preparation.

Table 1: Treatment-Emergent Adverse Reactions observed in at Least (>1%) of Patients using the Split-Dose Regimen and Day –Before Regimen **

Adverse Reaction	Study 1: Split-Dose Regimen		Study 2: Day-Before Regimen	
	PREPOPIK (N=305) n (% = n/N)	2L PEG+E* with 2 x 5-mg bisacodyl tablets (N=298) n (% = n/N)	PREPOPIK (N=296) n (% = n/N)	2L PEG+E* with 2 x 5-mg bisacodyl tablets (N=302) n (% = n/N)
Nausea	8 (2.6)	11 (3.7)	9 (3.0)	13 (4.3)
Headache	5 (1.6)	5 (1.7)	8 (2.7)	5 (1.7)
Vomiting	3 (1.0)	10 (3.4)	4 (1.4)	6 (2.0)

* 2L PEG + E = two liters polyethylene glycol plus electrolytes solution.

**abdominal bloating, distension, pain/cramping, and watery diarrhea not requiring an intervention were not collected

Electrolyte Abnormalities

In general, PREPOPIK was associated with numerically higher rates of abnormal electrolyte shifts on the day of colonoscopy compared to the preparation containing 2L of PEG + E plus two x 5-mg bisacodyl tablets (Table 2). These shifts were transient in nature and numerically similar between treatment arms at the Day 30 visit.

Table 2: Shifts from Normal Baseline to Outside the Normal Range at Day 7 and Day 30

Laboratory Parameter (direction of change)	Visit	Study 1: Split-Dose Regimen		Study 2: Day-Before Regimen	
		PREPOPIK	2L PEG+E with 2x 5 mg bisacodyl tablets	PREPOPIK	2L PEG+E with 2x 5 mg bisacodyl tablets
		n/N (%)		n/N (%)	
Potassium (low)	Day of Colonoscopy	19/260 (7.3)	11/268 (4.1)	13/274 (4.7)	13/271 (4.8)
	24-48 hours	3/302 (1.0)	2/294 (0.7)	3/287 (1.0)	5/292 (1.7)
	Day 7	11/285 (3.9)	8/279 (2.9)	6/276 (2.2)	14/278 (5.0)
Sodium (low)	Day 30	11/284 (3.9)	8/278 (2.9)	7/275 (2.5)	8/284 (2.8)
	Day of Colonoscopy	11/298 (3.7)	3/295 (1.0)	3/286 (1.0)	3/295 (1.0)
	24-48 hours	1/303 (0.3)	1/295 (0.3)	1/288 (0.3)	1/293 (0.3)
	Day 7	2/300 (0.7)	1/292 (0.3)	1/285 (0.4)	1/291 (0.3)

	Day 30	2/299(0.7)	3/291 (1.0)	1/284(0.4)	1/296 (0.3)
Chloride (low)	Day of Colonoscopy	11/301 (3.7)	1/298 (0.3)	3/287 (1.0)	0/297 (0.0)
	24-48 hours	1/303 (0.3)	0/295 (0.0)	2/288 (0.7)	0/293 (0.0)
	Day 7	1/303 (0.3)	3/295 (1.0)	0/285 (0.0)	0/293 (0.0)
	Day 30	2/302 (0.7)	3/294 (1.0)	0/285 (0.0)	0/298 (0.0)
Magnesium (high)	Day of Colonoscopy	34/294 (11.6)	0/294 (0.0)	25/288 (8.7)	1/289 (0.3)
	24-48 hours	0/303 (0.0)	0/295 (0.0)	0/288 (0.0)	0/293 (0.0)
	Day 7	0/297 (0.0)	1/291 (0.3)	1/286 (0.3)	1/285 (0.4)
	Day 30	1/296 (0.3)	2/290 (0.7)	0/286 (0.0)	0/290 (0.0)
Calcium (low)	Day of Colonoscopy	2/292 (0.7)	1/286 (0.3)	0/276 (0.0)	2/282 (0.7)
	24-48 hours	0/303 (0.0)	0/295 (0.0)	0/288 (0.0)	0/293 (0.0)
	Day 7	0/293 (0.0)	1/283 (0.4)	0/274 (0.0)	0/278 (0.0)
	Day 30	0/292 (0.0)	1/282 (0.4)	0/274 (0.0)	1 /283(0.4)
Creatinine (high)	Day of Colonoscopy	5/260 (1.9)	13/268 (4.9)	12/266 (4.5)	16/270 (5.9)
	24-48 hours	1/303 (0.3)	0/295 (0.0)	0/288 (0.0)	0/293 (0.0)
	Day 7	10/264 (0.4)	13/267 (4.8)	10/264 (3.8)	10/265 (3.8)
	Day 30	11/264 (4.2)	14/265(5.3)	18/264 (6.8)	10/272 (3.7)
eGFR (low)	Day of Colonoscopy	22/221 (10.0)	17/214 (7.9)	26/199 (13.1)	25/224 (11.2)
	24-48 hours	76/303 (25.1)	72/295 (24.4)	82/288 (28.5)	62/293 (21.2)
	Day 7	22/223 (10.0)	17/213 (8.0)	11/198 (5.6)	28/219 (12.8)
	Day 30	24/223(10.8)	21/211 (10.0)	21/199 (10.6)	24/224 (10.7)

6.2 Postmarketing Experience

The following foreign spontaneous reports have been identified during use of formulations similar to PREPOPIK. Because these events are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

Allergic reactions

Cases of hypersensitivity reactions including rash, urticaria, and purpura have been reported.

Electrolyte abnormalities

There have been reports of hypokalemia, hyponatremia and hypermagnesemia with the use of PREPOPIK for colon preparation prior to colonoscopy.

Gastrointestinal:

Abdominal pain, diarrhea, fecal incontinence, and proctalgia have been reported with the use of PREPOPIK for colon preparation prior to colonoscopy. There have been isolated reports of reversible aphthoid ileal ulcers. Ischemic colitis has been reported with the use of PREPOPIK for colon preparation prior to colonoscopy. However, a causal relationship between these ischemic colitis cases and the use of PREPOPIK has not been established.

Neurologic,

There have been reports of generalized tonic-clonic seizures associated with and without hyponatremia in epileptic patients.

7 DRUG INTERACTIONS

7.1 Drugs That May Increase Risks of Fluid and Electrolyte Abnormalities

Use caution when prescribing PREPOPIK for patients with conditions or who are using medications that increase the risk for fluid and electrolyte disturbances or may increase the risk of seizure, arrhythmias, and prolonged QT in the setting of fluid and electrolyte abnormalities. This includes patients receiving drugs which may be associated with hypokalemia (such as diuretics or corticosteroids, or drugs where hypokalemia is a particular risk, such as cardiac glycosides) or hyponatremia. Use caution when PREPOPIK is used in patients on nonsteroidal anti-inflammatory drugs (NSAIDs) or drugs known to induce Antidiuretic Hormone Secretion (SIADH), such as tricyclic antidepressants, selective serotonin re-uptake inhibitors, antipsychotic drugs and carbamazepine, as these drugs may increase the risk of water retention and/or electrolyte imbalance. Consider additional patient evaluations as appropriate. [see *Adverse Reactions (6.1, 6.2)*]

7.2 Potential for Altered Drug Absorption

Oral medication administered within one hour of the start of administration of PREPOPIK solution may be flushed from the GI tract and the medication may not be absorbed.

Tetracycline and fluoroquinolone antibiotics, iron, digoxin, chlorpromazine and penicillamine, should be taken at least 2 hours before and not less than 6 hours after administration of PREPOPIK to avoid chelation with magnesium.

7.3 Antibiotics

Prior or concomitant use of antibiotics with PREPOPIK may reduce efficacy of PREPOPIK as conversion of sodium picosulfate to its active metabolite BHPM is mediated by colonic bacteria.

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Pregnancy Category B

Reproduction studies with PREPOPIK have been performed in pregnant rats at oral doses up to 2000 mg/kg/day (about 1.2 times the recommended human dose based on the body surface area), and did not reveal any evidence of impaired fertility or harm to the fetus due to PREPOPIK. The reproduction study in rabbits was not adequate, as treatment-related mortalities were observed at all doses. A pre and postnatal development study in rats showed no evidence of any adverse effect on pre and postnatal development at oral doses up to 2000 mg/kg twice daily (about 1.2 times the recommended human dose based on the body surface area). There are, however, no adequate and well-controlled studies in pregnant women. Because animal reproduction studies are not always predictive of human response, PREPOPIK should be used during pregnancy only if clearly needed.

8.3 Nursing Mothers

It is not known whether this drug is excreted in human milk. Because many drugs are excreted in human milk, caution should be exercised when PREPOPIK is administered to a nursing woman.

8.4 Pediatric Use

The safety and effectiveness of PREPOPIK in pediatric patients has not been established.

8.5 Geriatric Use

In controlled clinical trials of PREPOPIK, 215 of 1201 (18%) patients were 65 years of age or older. The overall incidence of treatment-emergent adverse events was similar among patients ≥ 65 years of age (73%) and patients < 65 years of age (71%). Among all patients ≥ 65 years of age, the proportion of patients with successful colon cleansing was greater in the PREPOPIK group (81.1%) than in the comparator group (70.9%).

8.7 Renal Insufficiency

Patients with impaired renal function or patients taking concomitant medications that may affect renal function (such as diuretics, angiotensin converting enzyme inhibitors, angiotensin receptor blockers, or non-steroidal anti-inflammatory drugs) may be at increased risk for further renal injury. Advise these patients of the importance of adequate hydration before during and after the use of PREPOPIK. Consider performing baseline and post-colonoscopy laboratory tests (electrolytes, creatinine, and BUN) in these patients. In patients with severely reduced renal function (creatinine clearance < 30 mL/min), accumulation of magnesium in plasma may occur. The signs and symptoms of hypermagnesemia may include, but are not limited to, diminished or absent deep tendon reflexes, somnolence, hypocalcemia, hypotension, bradycardia, muscle, respiratory paralysis, complete heart block, and cardiac arrest.

10 OVERDOSAGE

The patient who has taken an overdose should be monitored carefully, and treated symptomatically for complications.

11 DESCRIPTION


PREPOPIK (sodium picosulfate, magnesium oxide and anhydrous citric acid) for oral solution is provided in two packets, the contents of each to be dissolved in 5 ounces of cold water and consumed.

Each packet contains 10 mg sodium picosulfate, 3.5 g magnesium oxide and 12 g anhydrous citric acid. The product also contains the following inactive ingredients, potassium hydrogen carbonate, saccharine sodium, and spray dried orange flavor which contains acacia gum, lactose, ascorbic acid and butylated hydroxyanisole. The following is a description of the three active ingredients:

Sodium picosulfate is a stimulant laxative.

Sodium picosulfate

- Chemical name: 4,4'-(2-pyridylmethylene) diphenyl bis(hydrogen sulfate) disodium salt, monohydrate
- Chemical formula: $C_{18}H_{13}NNa_2O_8S_2 \cdot H_2O$
- Molecular weight: 499.4
- Structural formula:



- Sodium picosulfate


Magnesium citrate, which is formed in solution by the combination of magnesium oxide and anhydrous citric acid, is an osmotic laxative.

Magnesium oxide

- Chemical name: Magnesium oxide
- Chemical formula: Mg O
- Molecular weight: 40.3
- Structural formula: Mg O

Anhydrous citric acid

- Chemical name: 2-hydroxypropane-1,2,3-tricarboxylic acid
- Chemical formula: $C_6H_8O_7$
- Molecular weight: 192.1
- Structural formula:



Anhydrous citric acid

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Sodium picosulfate is hydrolyzed by colonic bacteria to form an active metabolite: bis-(p-hydroxy-phenyl)-pyridyl-2-methane, BHPM, which acts directly on the colonic mucosa to stimulate colonic peristalsis.

Magnesium oxide and citric acid react to create magnesium citrate in solution, which is an osmotic agent that causes water to be retained within the gastrointestinal tract.

12.2 Pharmacodynamics

The stimulant laxative activity of sodium picosulfate together with the osmotic laxative activity of magnesium citrate produces a purgative effect which, when ingested with additional fluids, produces watery diarrhea.

12.3 Pharmacokinetics

Sodium picosulfate, which is a prodrug, is converted to its active metabolite, BHPM, by colonic bacteria. After administration of 2 packets of PREPOPIK separated by 6 hours, in 16 healthy volunteers, sodium picosulfate reached a mean C_{max} of 3.2 ng/mL at approximately 7 hours (T_{max}). After the first packet the corresponding values were 2.3 ng/mL at 2 hours. The terminal half-life of sodium picosulfate was 7.4 hours. The fraction of the absorbed sodium picosulfate dose excreted unchanged in urine was 0.19%.

Plasma levels of the free BHPM were low, with 13 out of 16 subjects studied having plasma BHPM concentrations below the lower limit of quantification (0.1 ng/mL). Urinary samples show that the majority of excreted BHPM was in the glucuronide-conjugated form. Magnesium oxide and citric acid react in water to create magnesium citrate. Baseline uncorrected magnesium concentration reached a maximum (C_{max}) of approximately 1.9 mEq/L, which occurred at 10 hours post initial packet administration (T_{max}). This represent an approximately 20% increase from the baseline.

Drug Interaction Studies

In an *in vitro* study using human liver microsomes, sodium picosulfate did not inhibit the major CYP enzymes (CYP 1A2, 2B6, 2C8, 2C9, 2C19, 2D6 and 3A4/5) evaluated. Based on an *in vitro* study using freshly isolated hepatocyte culture, sodium picosulfate is not an inducer of CYP1A2, CYP2B6 or CYP3A4/5.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Long-term studies in animals to evaluate carcinogenic potential or studies to evaluate mutagenic potential have not been performed with PREPOPIK. However, sodium picosulfate was not mutagenic in the Ames test, the mouse lymphoma assay and the mouse bone marrow micronucleus test.

In an oral fertility study in rats, PREPOPIK did not cause any significant adverse effect on male or female fertility parameters up to a maximum dose of 2000 mg/kg twice daily (about 1.2 times the recommended human dose based on the body surface area).

14 CLINICAL STUDIES

The colon cleansing efficacy of PREPOPIK was evaluated for non-inferiority against a comparator in two randomized, investigator-blinded, active-controlled, multicenter US trials in patients scheduled to have an elective colonoscopy. In all, 1195 adult patients were included in the primary efficacy analysis: 601 from Study 1, and 594 from Study 2. Patients ranged in age from 18 to 80 years (mean age 56 years); 61% were female and 39% male. Self-identified race was distributed as follows: 90% White, 10% Black, and less than 1% other. Of these, 3% self-identified their ethnicity as Hispanic or Latino.

Patients randomized to Prepopik in the two studies were treated with one of two dosing regimens:

- In Study 1, PREPOPIK was given by “Split-Dose” (evening before and day of) dosing, where the first packet was taken the evening before the colonoscopy (between 5:00 and 9:00 PM), followed by five (5) 8-ounce glasses of clear liquid, and the second packet was taken the morning of the colonoscopy (at least 5 hours prior to but no more than 9 hours prior to colonoscopy), followed by three (3) 8-ounce glasses of clear liquid.
- In Study 2, PREPOPIK was given by “Day-Before” (afternoon/evening before only) dosing, where both packets were taken separately on the day before the colonoscopy, with the first packet taken in the afternoon (between 4:00 and 6:00 PM), followed by five (5) 8-ounce glasses of clear liquid, and the second packet taken in the late evening (approximately 6 hours later, between 10:00 PM and 12:00 AM), followed by three (3) 8-ounce glasses of clear liquid.

The comparator was a preparation containing two liters of polyethylene glycol plus electrolytes solution (PEG + E) and two 5-mg bisacodyl tablets, administered the day before the procedure. All patients in both the Prepopik and comparator groups were limited to a clear liquid diet on the day before the procedure (24 hours before).

The primary efficacy endpoint was the proportion of patients with successful colon cleansing, as assessed by blinded colonoscopists using the Aronchick Scale. The Aronchick scale is a tool used to assess overall colon cleansing. Successful colon cleansing was defined as bowel preparations with >90% of the mucosa seen and mostly liquid stool that were graded excellent (minimal suctioning needed for adequate visualization) or good (significant suctioning needed for adequate visualization) by the colonoscopist.

In both studies, PREPOPIK was non-inferior to the comparator. In addition, PREPOPIK provided by Split-Dose dosing met the pre-specified criteria for superiority to the comparator for colon cleansing in Study 1. The comparator in that study was administered entirely on the day prior to colonoscopy. See Tables 3 and 4 below.

Table 3: Proportion of Patients with Successful Colon Cleansing in Study 1 Split –Dose Regimen

PREPOPIK Split-Dose Regimen	2L PEG+E* with 2 x 5-mg bisacodyl tablets	Difference between treatment groups	
% (n/N)	% (n/N)	Difference	95% CI
84.2% (256/304)	74.4% (221/297)	9.8%	(3.4%, 16.2%) [†]

* 2L PEG + E = two liters polyethylene glycol plus electrolytes solution.

[†] Non-inferior and superior 2L PEG+E with 2 x 5-mg bisacodyl tablets

Table 4: Proportion of Patients with Successful Colon Cleansing in Study 2 Day-Before Regimen

PREPOPIK Day-Before Regimen	2L PEG+E* with 2 x 5-mg bisacodyl tablets	Difference between treatment groups	
% (n/N)	% (n/N)	Difference	95% CI
83.0% (244/294)	79.7% (239/300)	3.3%	(-2.9%, 9.6%) [‡]

* 2L PEG + E = two liters polyethylene glycol plus electrolytes solution.

[‡]. Non-inferior

16 HOW SUPPLIED/STORAGE AND HANDLING

How Supplied

PREPOPIK is supplied in a carton containing 2 packets, each holding 16.1 grams of powder for oral solution, along with a pre-marked dosing cup. Each packet contains 10 mg sodium picosulfate, 3.5 g magnesium oxide and 12 g anhydrous citric acid. The excipients include potassium hydrogen carbonate, sodium saccharin, spray dried orange flavor which contains acacia gum, lactose, ascorbic acid, and butylated hydroxyanisole.

Storage

Store at 25°C (77°F). Excursions permitted at 15°C to 30°C (59°F to 86°F) [See USP Controlled Room Temperature].

NDC# 55566-9300-2 Kit, 2 packets and cup

17 PATIENT COUNSELING INFORMATION

See FDA-approved patient labeling (Medication Guide).

- Ask patients to let you know if they have trouble swallowing or are prone to regurgitation or aspiration.
- Tell patients not to take other laxatives while they are taking PREPOPIK.
- Tell patients that if they experience severe bloating, distention or abdominal pain following the first packet of PREPOPIK, delay the second administration until the symptoms resolve.
- Instruct patients to contact their healthcare provider if they develop signs and symptoms of dehydration.
- Not for Direct Ingestion: Each packet must be dissolved in 5 ounces of cold water and administered at separate times according to the dosing regimen. Ingestion of additional water is important to patient tolerance. Direct ingestion of the undissolved powder may increase the risk of nausea, vomiting, dehydration, and electrolyte disturbances. Inform patients that oral medication administered within one hour of the start of administration of PREPOPIK solution may not be absorbed completely.

Manufactured by:

Ferring Pharmaceuticals (China) Co., Ltd.
No. 6 HuiLing Lu (Ferring Road)
National Health Technology Park
Zhongshan City, Guangdong Province, CHINA

Manufactured for:

Ferring Pharmaceuticals Inc.
Parsippany, N.J. 07054

Medication Guide

PREPOPIK (prep-ō-pik)

(sodium picosulfate, magnesium oxide and anhydrous citric acid) for oral solution

Read this Medication Guide instructions before you start taking PREPOPIK and each time you get a refill. There may be new information. This information does not take the place of talking with your healthcare provider about your medical condition or your treatment.

What is the most important information I should know about PREPOPIK?

PREPOPIK and other bowel preparations can cause serious side effects, including:

Serious loss of body fluid (dehydration) and changes in blood salts (electrolytes) in your blood. These changes can cause:

- abnormal heartbeats that can cause death
- seizures. This can happen even if you have never had a seizure.
- kidney problems

Your chance of having fluid loss and changes in blood salts with PREPOPIK is higher if you:

- have heart problems
- have kidney problems
- take water pills or non-steroidal anti-inflammatory drugs (NSAIDs)

Tell your healthcare provider right away if you have any of these symptoms of a loss of too much body fluid (dehydration) while taking PREPOPIK:

- vomiting that prevents you from keeping down the additional prescribed amounts of clear liquids that you must drink after taking your PREPOPIK
- dizziness
- urinating less often than normal
- headache

See “What are the possible side effects of PREPOPIK?” for more information about side effects.

What is PREPOPIK?

PREPOPIK is a prescription medicine used by adults to clean the colon before a colonoscopy. Prepopik cleans your colon by causing you to have diarrhea. Cleaning your colon helps your healthcare provider see the inside of your colon more clearly during your colonoscopy.

It is not known if PREPOPIK is safe and effective in children.

Who should not take PREPOPIK?

Do not take PREPOPIK if your healthcare provider has told you that you have:

- serious kidney problems
- a blockage in your intestine (bowel obstruction)
- an opening in the wall of your stomach or intestines (bowel perforation)
- a very dilated intestine (toxic megacolon)
- problems with the emptying of food and fluid from your stomach (gastric retention)
- an allergy to any of the ingredients in PREPOPIK. See the end of this leaflet for a complete list of ingredients in PREPOPIK.

What should I tell my healthcare provider before taking PREPOPIK?

Before you take PREPOPIK, tell your healthcare provider if you:

- have heart problems
- have stomach or bowel problems
- have ulcerative colitis
- have problems with swallowing or gastric reflux
- are withdrawing from drinking alcohol and benzodiazepines
- have kidney problems
- have low blood salt (sodium) level
- any other medical conditions
- are pregnant. It is not known if Prepopik will harm your unborn baby. Talk to your provider if you are pregnant or plan to become pregnant.
- are breastfeeding or plan to breastfeed. It is not known if Prepopik passes into your breast milk. You and your healthcare provider should decide if you will take Prepopik while breastfeeding.

Tell your healthcare provider about all the medicines you take, including prescription and non-prescription medicines, vitamins, and herbal supplements.

Prepopik may affect how other medicines work. Medicines taken by mouth may not be absorbed properly when taken within 1 hour before the start of PREPOPIK.

Especially tell your healthcare provider if you take:

- medicines for blood pressure or heart problems
- medicines for kidney problems
- medicines for seizures
- water pills (diuretics)
- nonsteroidal anti-inflammatory medicines (pain medicines)
- medicines for depression or mental health problems
- laxatives
- the following medicines should be taken at least 2 hours before starting PREPOPIK and not less than 6 hours after taking PREPOPIK:
 - tetracycline
 - fluoroquinolone antibiotics
 - iron
 - digoxin (Lanoxin)
 - chlorpromazine
 - penicillamine (Cuprimine, Depen)

Ask your healthcare provider or pharmacist for a list of these medicines if you are not sure if you are taking the medicines listed above.

Know the medicines you take. Keep a list of them to show your healthcare provider and pharmacist when you get a new medicine.

How should I take PREPOPIK?

See the Instructions for Use on the outer product carton for dosing. You must read, understand, and follow these instructions to take PREPOPIK the right way.

- **Take PREPOPIK exactly as your healthcare provider tells you to take it.** Your healthcare provider will prescribe the Split-Dosing option or the Day Before Dosing option, depending on colonoscopy scheduling, distance traveled, and other personal circumstances.
- A complete preparation requires 2 packets of PREPOPIK for oral solution taken separately, each followed by additional fluids.
- It is important for you to drink the additional prescribed amount of clear liquids after taking PREPOPIK to prevent fluid loss (dehydration).

- Examples of clear liquids include water, clear broth, apple juice, white cranberry juice, white grape juice, and ginger ale, plain jello (not red or purple) and frozen juice bars(not purple or red).
- Do not eat solid foods or drink milk while taking PREPOPIK
- Drink clear liquids until your colonoscopy.
- Do not take other laxatives while taking PREPOPIK.
- Stop drinking PREPOPIK temporarily or allow for longer time between each dose if you have bloating, distension, or stomach (abdominal) pain until your symptoms improve.
- Stop taking PREPOPIK and, and call your healthcare provider right away if you develop hives or rash after you take your first packet of PREPOPIK. These may be signs of an allergic reaction.

See the Instructions for Use on the outer product carton for dosing. You must read, understand, and follow these instructions to take PREPOPIK the right way.

1) Split-Dose (evening-before and day of the procedure) Dosing

Take your first packet of PREPOPIK the night before your colonoscopy, and take your second dose the next day, in the morning before your colonoscopy.

On the day before your colonoscopy procedure – 1 packet:

- Dissolve 1 packet of powder in 5 ounces of cold water in the evening, followed by five 8-ounce drinks (upper line on the dosing cup) of clear liquids before bed.

On the day of the colonoscopy procedure – 1 packet:

- Dissolve 1 packet of powder in 5 ounces of cold water in the morning (5 hours before the colonoscopy), followed by at least three 8-ounce drinks of clear liquids before the colonoscopy.

You may continue to drink clear liquids until 2 hours before the time of the colonoscopy.

2) Day-Before (afternoon and evening-before the procedure) Dosing

Take your first PREPOPIK packet in the afternoon or early evening and take your second packet 6 hours later, the night before the colonoscopy.

On the day before the colonoscopy procedure – 2 packets:

- Dissolve 1 packet of powder in 5 ounces of cold water in the afternoon or early evening, followed by five 8-ounce drinks (upper line on the dosing cup) of clear liquids before the next dose.
- Dissolve 1 packet of powder in 5 ounces of cold water in the late evening (followed by three 8-ounce drinks (upper line on the dosing cup) of clear liquids before bed.

You may continue to drink clear liquids until 2 hours before the time of the colonoscopy.

What are the possible side effects of PREPOPIK?

PREPOPIK can cause serious side effects, including:

See **“What is the most important information I should know about PREPOPIK?”**

- **changes in certain blood tests.** Your healthcare provider may do blood tests after you take PREPOPIK to check your blood for changes. Tell your healthcare provider if you have any symptoms of too much fluid loss, including:
 - Vomiting
 - Nausea
 - Bloating
 - Dizziness
 - Stomach(abdominal)cramping

- Urinate less than usual
- Trouble drinking clear liquids
- Troubles swallowing
- Seizures
- Heart problems (arrhythmia). PREPOPIK may cause irregular heartbeats
- Ulcers of the bowel or bowel problems (ischemic colitis). Tell your healthcare provider right away if you have severe stomach (abdominal) pain or rectal bleeding. These may be symptoms of decreased blood flow to the intestine.

The most common side effects of PREPOPIK include:

- nausea
- headache
- vomiting

Tell your healthcare provider if you have any side effect that bothers you or that does not go away.

These are not all the possible side effects of PREPOPIK. For more information, ask your healthcare provider or pharmacist.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

How should I store PREPOPIK?

- Store PREPOPIK at room temperature, between 68 to 77°F (20 to 25°C).

Keep PREPOPIK and all medicines out of the reach of children.

General information about the safe and effective use of PREPOPIK.

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. Do not use PREPOPIK for a condition for which it was not prescribed. Do not give PREPOPIK to other people, even if they are going to have the same procedure you are. It may harm them.

This Medication Guide summarizes the most important information about PREPOPIK. If you would like more information, talk with your healthcare provider. You can also ask your pharmacist or healthcare provider for information that is written for healthcare professionals.

For more information, go to www.ferring.com or call 1-888-337-7464.

What are the ingredients in PREPOPIK?

Active ingredients: sodium picosulfate, magnesium oxide and anhydrous citric acid

Inactive ingredients: potassium hydrogen carbonate, saccharin sodium, spray dried orange flavor which contains acacia gum, lactose, ascorbic acid and butylated hydroxyanisole

This Medication Guide has been approved by the U.S. Food and Drug Administration.

Ferring Pharmaceuticals Inc.
Parsippany, NJ 07054, USA

Issued July/2012