RYAN CRENSHAW, M.D. Instructions for Colonoscopy with NuLytely

Please read this packet, in its entirety, at least 2 weeks prior to your procedure. If you fail to follow the instructions and the procedure has to be cancelled, the cancellation fee will be charged.

If you need to cancel your procedure, please let us know at least <u>5 business days</u> prior to the procedure. If you fail to do so, you will be charged a \$250.00 cancellation fee.

You will receive a confirmation call from our office staff at least <u>5 days</u> prior to your scheduled procedure(s). All procedure(s) must be confirmed in order to remain on the schedule. If you do not receive a call from our office, please contact us at 703-444-4799.

If you take blood thinners such as Aspirin, Plavix, Xarelto or Coumadin, Dr. Crenshaw may recommend for you to hold these medications anywhere from 2-8 days prior to your procedure, depending on what agent you are taking.

Patients on Coumadin (Warfarin): If approved by the prescribing physician (i.e. cardiologist, neurologist or primary care provider), you will be asked to stop your Coumadin 5 days prior to your procedure(s). You will also be asked to obtain a PT/INR, PTT blood test the day prior to your procedure(s). If you did not receive an order for this blood test, please contact our office at (703) 444-4799. Dr. Crenshaw will instruct you regarding the date to restart Coumadin (Warfarin) on the day of your procedure(s).

Patients on Plavix (Clopidogrel): If approved by the prescribing physician (i.e. cardiologist, neurologist or primary care provider), you will be asked to stop the Plavix (clopidrogel) 7 days prior to your procedure(s). Dr. Crenshaw will instruct you regarding the date to restart Plavix (Clopidogrel) on the day of your procedure(s).

Patients taking Aspirin: Please make sure one of the two lines is checked off below. If not, please contact our office at (703) 444-4799.

Please continue to t	ake Aspirin	o. 81 mg or 325 mg	. daily, including	g the day	v of the colonoscopy
i icase continue to i	and Aspirii	1, UI INS OF JAJ INS	, auny, menann	S LIIC GG	y of the colonoscopy

__ Please stop taking Aspirin, 81 mg or 325 mg, 8 days prior to the colonoscopy,

Any patient stopping Aspirin, Plavix, Coumadin or any other blood thinner should contact the prescribing doctor (primary care physician, or cardiologist) to confirm that it is acceptable to stop this medication(s) for the recommended period of time. You may take Tylenol if needed. Please do not take any other medication or products that can thin the blood, such as Ibuprofen, Motrin, Advil, Aleve, Vitamin E and/or Garlic pills 8 days prior to the procedure(s).

It is highly recommended that you take your medication for heart disease, high blood pressure and asthma every day, including the day of your procedure. If you take any medication(s) around the time that you are taking a dose of the laxative to prepare for your colonoscopy, please take the laxative first, then your medication. When taking medication(s) on the day of your procedure, the medication(s) must be taken at least 4 hours prior to your procedure time with water. At 4 hours prior to procedure time, you should stop consuming liquids, stop taking medications and not take anything orally until your procedure has been completed. All other medications should be brought to the hospital to be taken after your procedure.

If you are taking medications for diabetes, consult with the medical provider that is managing your diabetes to inform him/her that you are being asked to change your diet in preparation for colonoscopy. Please ask this provider how you should change your diabetes medication regimen to reduce the risk of your blood sugar becoming too low or too high during your preparation for the procedure.

You will need someone to drive you home from the hospital, or surgical center, after your procedure. You should not drive until the next day.

Preparation Instructions for NuLytely

(Polyethylene glycol 3350, 420 g, Sodium bicarbonate 5.72 g, Sodium Chloride 11.2 g and Potassium Chloride 1.4 g)

Caution: This preparation must be used with caution and may be contraindicated in patients with the following conditions: Kidney failure or compromise in kidney function, uncorrected electrolyte abnormalities, heart arrhythmias, history of seizure, impaired gag reflex, gastric retention, gastrointestinal obstruction and/or severe ulcerative colitis. If you have any of these conditions, please call the office to confirm this preparation is appropriate for you.

Five days prior to the procedure: Do NOT eat food containing seeds, corn, nuts, black pepper, lettuce, raw vegetables, fruits with seeds or skin as they can be difficult to lavage from the colon. Please do not take fish oil, krill oil, lemon oil or any supplements or foods that contain a significant amount of oil such as potato chips.

The day prior to your procedure, you will be on a clear liquid diet (no solid food, except for Jello) for the entire day beginning with your breakfast meal. On the day before your colonoscopy, the more clear liquids you consume, the better for your preparation. In addition, adequate hydration will reduce the risk of developing headaches, lightheadedness and dizziness which can occur during the preparation.

It is also important to vary the liquids you are consuming. Do NOT restrict yourself to drinking water only. Please make sure you consume liquids with salt such as soup broth. And liquids with carbohydrates/sugars such as apple juice. By varying the liquids consumed, you reduce the risk of developing electrolyte abnormalities such as a low sodium. Please avoid any food or beverage product(s) which contain red or purple coloring.

MENU FOR CLEAR LIQUID DIET

Breakfast Lunch <u>Dinner</u> White Cranberry Juice Chicken broth Chicken broth White Grape Juice Gelatin dessert Apple Juice Tea/coffee (no milk) Sprite, 7up, Ginger Ale Gelatin dessert Gatorade or similar sports Fruit-flavored ice Sprite, 7up, Ginger Ale drink Tea/coffee (no milk) Tea/coffee (no milk)

Instructions continue on the next page

NULYTELY MEDICATION FOLLOW THE INSTRUCTIONS WRITTEN BELOW

FIRST DOSE: One day prior to your procedure, at 6:00 pm, you will need to consume 2 liters (1/2 gallon) of NuLytely.

You should try to consume an 8 ounce glass every 10 minutes to be completed by 8 pm. You can then consume clear liquids until 2 hours prior to bedtime from the menu selections listed above.

*Patients may suck on sugar-free menthol candy to alter/enhance the taste if desired.

SECOND DOSE: Day of the procedure: At 7 hours prior to your procedure time, please begin consuming the remaining 2 liters (1/2 gallon) of NuLytely. Try to consume an 8 ounce glass every 10 minutes to be completed by 5 hours prior to the scheduled procedure time.

*Patients may suck on sugar-free menthol candy to alter/enhance the taste if desired.

After completing the NuLytely solution, you can consume up to 24 ounces of clear liquids over the next 60 minutes. You must be done with the clear liquids 4 hours prior to your scheduled procedure time.

Please note that consuming any liquids or medication(s) too close to your procedure may result in delaying or cancelling your procedure. Do not consume any liquids or medication(s) by mouth for 4 hours prior to your procedure time.

On average, your bowel movements should be clear (clear = yellow or white liquid without solid or granular material) 3 hours after you finish drinking the solution. However, delays in the onset of bowel movements and/or becoming clear can occur several hours after completing the solution. If you are not clear at 2 hours after taking the second dose on the day of your procedure, please follow the instructions written below.

If you are unable to complete and/or tolerate the preparation for colonoscopy, please follow these instructions:

Purchase the following (no prescription necessary) and begin this preparation 1/2 hour after the last glass of solution taken.

- One bottle of Magnesium Citrate
- One bottle of fleet enema.

Drink one bottle of Magnesium Citrate. Wait 2 hours. If your bowel movements are not clear or you could not tolerate the Magnesium Citrate, then proceed with Fleet Enema as follows:

Instuctions continue on the next page

- 1. Apply one Fleet Enema per rectum and wait 30 minutes.
- 2. If your bowel movements are still not clear, fill the same enema bottle with warm water from the faucet. Then administer a warm water enema every 30 minutes until you have bowel movements which consist of clear yellow or clear white liquid. Do NOT exceed more than 4 enemas.

If you still encounter significant difficulties with your preparation, please contact our office at (703) 444-4799. If you are forwarded to voicemail, follow directions to contact the doctor on call.

RYAN P. CRENSHAW, M.D.

21135 WHITFIELD PLACE, SUITE 102, STERLING, VA 20165 (703) 444-4799

OPERATIVE REQUEST/CONSENT

1.	procedure(s) along with surgice	al assistants selected by herforation, and need for	enshaw (the "Practitioner") to perform: colonoscopy, possible biopsy, risk or surgery has been explained. The Practic "Patient'):	f drug allergy, over					
	Please print your name:								
2.	2. It has been clearly explained to me that during the course of this operation some other conditions that have not been expected may present themselves. I recognize, that if such conditions are discovered it will be necessary to more than that which was specified in paragraph #1 above. I therefore authorize and request that the above named Practitioner and his surgical assistants perform such surgical procedures which in their best professions judgment will be effective in their attempt to heal and/or diagnose. This includes, but is not limited to, patholog and radiology. I further authorize Anesthesiologist to administer whatever anesthesia they feel is indicated an authorize the use of blood transfusion(s) when attending personnel feel such is required.								
3.	I fully understand that this ope is guaranteed.	ration, like any operation	, is accompanied by some degree of ris	k and that no cure					
4.	risks involved and whatever ot	her choices are available en given the opportunit	of the procedure(s) listed under paragr to me (or the patient), if any, have bee y to ask any questions that I may ha factorily.	n explained to me					
5.	I am aware of the "Cancellation not provided at least 5 busines		that I will be held responsible for a \$250 duled date for procedure.	0.00 fee if notice is					
6.	· · · · · · · · · · · · · · · · · · ·		h both verbal and written instructions g your procedure), you will be charged th						
7.	procedure(s), as well as obtaini	ng the necessary referral	surance provider to check coverage f s. It is also the patient's responsibility t se, the patient will be held responsible f	o notify our office					
Sig	nature of Patient	Date	Signature of Witness	Date					
igr	nature of Parent/Guardian	Date							
l ł	• • • •	•	EMENT posed procedure to the patient, and/or his procedure, and any alternative.	· Relative/guardian,					
	_	Signature of Physicia	n Date						
		B = f=	n / 10 11	. 1					

Page 5 of 5

Patient Copy: Nulytely

RYAN P. CRENSHAW, M.D.21135 WHITFIELD PLACE, SUITE 102, STERLING, VA 20165 (703) 444-4799

OPERATIVE REQUEST/CONSENT

	not provided at least <u>5 busine</u> If your procedure is cancelled	*		250.00 fee if not	ce is
·5.	by the Practitioner. I have be explanation and my questions. I am aware of the "Cancellation"	een given the opportunity to have been answered satisfactors. On Policy" and understand that	nt I will be held responsible for a \$	have regarding	that
3.	is guaranteed. The nature of my (or the patie	nt's) condition, the nature of	accompanied by some degree of the procedure(s) listed under par	agraph #1 above	, the
2.	It has been clearly explained been expected may present to do more than that which was named Practitioner and his sigudgment will be effective in and radiology. I further author	hemselves. I recognize, that i specified in paragraph #1 ab urgical assistants perform suc their attempt to heal and/or o prize Anesthesiologist to adm	of this operation some other corf such conditions are discovered ove. I therefore authorize and rech surgical procedures which in the diagnose. This includes, but is not hinister whatever anesthesia they personnel feel such is required.	it will be necessa quest that the ab heir best professi limited to, patho	ary to bove siona olog
2.	• •	to me that during the course	of this operation some other cor		