## Medications Safe to Take During Pregnancy

**Cold/Flu/Congestion/Cough**: Sudafed (after 1<sup>st</sup> trimester), Actifed (after 1<sup>st</sup> trimester), Mucinex, Robitussin, Delsym, Vicks Dayquil \***Avoid Pseudoephedrine and Phenylephrine in 1<sup>st</sup> trimester and with high blood pressure.** 

Sore Throat: Cough lozenges, warm saline gargle (1tsp.salt in 1 quart of warm water)

Allergy Symptoms: Benadryl, Claritin, Zyrtec, Alavert, chlorpheniramine, Rhinocort

Rash: Cortaid topical, Benadryl cream, hydrocortisone cream, Caladryl

**Constipation**: Metamucil, Dulcolax, FiberCon, Colace, Milk of Magnesia, Miralax, Senekot, Citrucil

**Fever**: Tylenol (call if > 101.5 after taking Tylenol)

Headache: Tylenol

Pain and Discomfort: Tylenol

Diarrhea: Immodium A-D - Limit to 24 hours of use (avoid in 1st trimester)

Heartburn: Tums, Maalox, Mylanta, Zantac, Pepcid

Nausea: Emetrol, Unisom, Mylanta, Vitamin B6, peppermint

Hemorrhoids: Preparation H, Anusol, Tucks Pads

Yeast Infection: Clotrimazole, Monistat (miconazole)

Sleep: Unisom, Benadryl

Gas: simethicone

\*Please note: No drug can be considered 100% safe.