































## Prevention







Goal 1: Reduce the incidence and mortality of tobacco-related cancers									
Indicator	Measure	Baseline	January 2017	October 2017	March 2019	June 2020	2020 Target	Progress towards target	Target Met
1.1	Percentage of adults who are current smokers (age adjusted to the year 2000 standard population)	16.5% BRFSS 2014	14.2% BRFSS 2015	15.0% BRFSS 2016	14.8% BRFSS 2017	15.4% BRFSS 2018	12.0% HP2020		
1.2	Percentage of adolescents in grades 9 through 12 who used cigarettes, chewing tobacco, snuff, or cigars in the past 30 days	17.8% YRBS 2013 (rev)	17.4% YRBS 2015		12.5% YRBS 2017	8.2% YRBS 2019	14.2% CCAI (20%)		
1.3	Percentage of adult males aged 18+ who are current users of smokeless tobacco products such as chewing tobacco, snuff, and snus (age adjusted to the year 2000 standard population)	9.4% BRFSS 2014	9.8% BRFSS 2015	11.8% BRFSS 2016	10.2% BRFSS 2017	8.4% BRFSS 2018	7.5% CCAI (20%)		
Goal 2: Increase access to healthy food options and opportunities for physical activity									
Indicator	Measure	Baseline	January 2017	October 2017	March 2019	June 2020	2020 Target	Progress towards target	Target Met
2.1	Percentage of adults aged 18+ who engage in the recommended level of weekly physical activity (age adjusted to the year 2000 standard population)	20.5% BRFSS 2013 (rev)	21.4% BRFSS 2015		23.7% BRFSS 2017		24.6% CCAI (20%)		
2.2	Percentage of adults aged 20+ who are at a healthy weight (BMI >= 18.5 and <= 25.0; age adjusted to the year 2000 standard population)	32.5% BRFSS 2014	32.5% BRFSS 2015	33.1% BRFSS 2016	31.2% BRFSS 2017	33.0% BRFSS 2018	35.8% CCAI (10%)		
2.3	Percentage of adolescents in grades 9 through 12 who meet physical activity guidelines for aerobic physical activity	27.9% YRBS 2013	29.6% YRBS 2015		23.7% YRBS 2017	22.2% YRBS 2019	31.6% HP2020		

Goal 3: Increase protective behaviors from sun and other ultraviolet radiation exposure									
Indicator	Measure	Baseline	January 2017	October 2017	March 2019	June 2020	2020 Target	Progress towards target	Target Met
3.1	Percentage of adolescents in grades 9 through 12 who report using artificial sources of ultraviolet light for tanning	12.3% YRBS 2013 (rev)	9.4% YRBS 2015		5.9% YRBS 2017	5.1% YRBS 2019	9.8% CCAI (20%)		✓
3.2	Percentage of adults aged 18+ who report using artificial sources of ultraviolet light for tanning (age adjusted to the year 2000 standard population)	5.6% BRFSS 2014		3.3% BRFSS 2016			4.5% CCAI (20%)		✓
3.3	Percentage of adults aged 18+ who reported having at least one sunburn in the past 12 months (age adjusted to the 2000 US Standard Population)	49.2% BRFSS 2018 (rev)					44.3% CCAI (10%) (rev)		
Goal 4: Increase the vaccination rate for vaccines shown to reduce the risk of cancer									
Indicator	Measure	Baseline	January 2017	Oct 2017/ Jan 2018	March 2019	June 2020	2020 Target	Progress towards target	Target Met
4.1	Percentage of adolescent females aged 13-17 years who completed 3 doses of the HPV vaccine, or 2 doses 6 months apart if 1st dose before age 15	31.1% IRIS 2014	35.4% IRIS 2015	38.8% IRIS 2016	41.5% IRIS 2017	43.8% IRIS 2018 45.9% IRIS 2019	80.0% HP2020		■
4.2	Percentage of adolescent males aged 13-17 years who completed 3 doses of the HPV vaccine, or 2 doses 6 months apart if 1st dose before age 15	15.8% IRIS 2014	22.1% IRIS 2015	27.7% IRIS 2016	32.9% IRIS 2017	37.0% IRIS 2018 41.0% IRIS 2019	80.0% HP2020		■
4.3	Percentage of newborns receiving hepatitis B vaccine (Hepatitis B vaccine administered from birth through age 3 days)	83.3% IRIS 2014	80.2% IRIS 2015	80.2% IRIS 2016	78.4% IRIS 2017	74.4% IRIS 2018 73.3% IRIS 2019	85.0% HP2020		■










Goal 5: Reduce cancer risk related to environmental carcinogens									
Indicator	Measure	Baseline	January 2017	October 2017	March 2019	June 2020	2020 Target	Progress towards target	Target Met
5.1	Percentage of adults living in households ever been tested for radon (age adjusted to the year 2000 standard population)	20.7% BRFSS 2014		19.8% BRFSS 2016		23.4% BRFSS 2018	24.8% CCAI (20%)		

## Early Detection and Screening

Goal 6: Reduce breast cancer deaths and rate of late stage diagnosis through screening and early detection									
Indicator	Measure	Baseline	January 2017	Oct 2017/ Jan 2018	March 2019	June 2020	2020 Target	Progress towards target	Target Met
6.1	Percentage of women aged 50 to 74 who had a mammogram within the past two years (age adjusted to the year 2000 standard population)	68.9% BRFSS 2014		64.3% BRFSS 2016		67.8% BRFSS 2018	81.1% HP2020		
6.2	Age-adjusted rate per 100,000 females of breast cancer diagnoses at late stage (regional and distant)	42.8 CDRI 2013 (rev)	46.1 CDRI 2014	40.5 CDRI 2015	43.3 CDRI 2016 * Stage Change	47.0 CDRI 2017	38.5 CCAI (10%)		
6.3	Age-adjusted mortality rate, female breast cancer	20.7 BVRHS 2014	22.3 BVRHS 2015	21.4 BVRHS 2016	21.6 BVRHS 2017	22.1 BVRHS 2018	18.6 CCAI (10%)		
Goal 7: Reduce deaths and numbers of new cases of cervical cancer through screening and early detection									
Indicator	Measure	Baseline	January 2017	Oct 2017/ Jan 2018	March 2019	June 2020	2020 Target	Progress towards target	Target Met
7.1	Percentage of women aged 21-65 who have had a Pap test within the past three years (age adjusted to the year 2000 standard population)	76.3% BRFSS 2014		73.0% BRFSS 2016		70.4% BRFSS 2018	93.0% HP2020		
7.2	Age-adjusted rate per 100,000 females of invasive cervical cancer diagnoses	5.2 CDRI 2013	6.3 CDRI 2014	5.7 CDRI 2015	7.8 CDRI 2016	7.4 CDRI 2017	4.7 CCAI (10%)		
7.3	Age-adjusted cervical cancer mortality rate per 100,000 females	2.0 BVRHS 2014	1.8 BVRHS 2015	2.1 BVRHS 2016	1.5 BVRHS 2017	1.6 BVRHS 2018	1.6 CCAI (20%)		


Goal 8: Reduce the numbers of deaths and new cases of colorectal cancers through screening and early detection									
Indicator	Measure	Baseline	January 2017	Oct 2017/ Jan 2018	March 2019	June 2020	2020 Target	Progress towards target	Target Met
8.1	Percentage of adults aged 50-75 who reported receiving a colorectal cancer screening based on the most recent guidelines, which include a blood stool test in the past year, sigmoidoscopy in the past 5 years and blood stool test in the past 3 years, or a colonoscopy in the past 10 years. (age adjusted to the year 2000 standard population)	60.9% BRFSS 2014		61.9% BRFSS 2016		66.2% BRFSS 2018	80.0% NCCRT		
8.2	Age-adjusted rate per 100,000 of invasive colorectal cancer incidence	35.8 CDRI 2013 (rev)	36.1 CDRI 2014	36.2 CDRI 2015	35.1 CDRI 2016	33.5 CDRI 2017	32.2 CCAI (10%)		
8.3	Age-adjusted mortality rate, colorectal cancer	12.9 BVRHS 2014	12.3 BVRHS 2015	13.2 BVRHS 2016	13.1 BVRHS 2017	13.4 BVRHS 2018	11.6 CCAI (10%)		
Goal 9: Monitor the development and implementation of screening and early detection methods for other cancers									
	There are no measures established for this goal. Refer to the Idaho Comprehensive Cancer Strategic Plan for objectives and strategic actions.								

## Treatment

Goal 10: Increase timely access to quality cancer diagnostic and treatment services for all Idahoans									
Indicator	Measure	Baseline	January 2017	October 2017	March 2019	June 2020	2020 Target	Progress towards target	Target Met
10.1	Percentage of Idaho adults aged 18-64 with health care coverage (age adjusted to the year 2000 standard population)	79.3% BRFSS 2014	82.2% BRFSS 2015	80.6% BRFSS 2016	80.1% BRFSS 2017	80.1% BRFSS 2018	95.2% CCAI (20%)		
10.2	Percentage of Idahoans who could not see a doctor due to cost sometime in past year (age adjusted to the year 2000 standard population)	16.4% BRFSS 2014	14.3% BRFSS 2015	14.7% BRFSS 2016	14.1% BRFSS 2017	15.8% BRFSS 2018	13.1% CCAI (20%)		
10.3	5-year relative survival ratio, adjusted for age and primary site mix (NAACCR cancer survival index)	63.6 CDRI 05-11	63.9 CDRI 06-12	64.4 CDRI 07-13	64.2 CDRI 08-14	64.7 CDRI 09-15	65.6 CCAI (Best states)		
Goal 11: Increase opportunities to access and participate in cancer treatment clinical trials									
Indicator	Measure	Baseline	January 2017	October 2017	March 2019	June 2020	2020 Target	Progress towards target	Target Met
11.1	Percentage of cancer patients who enroll in treatment-related clinical trials	20.5% Ages 0-19		23.3% Ages 0-19	12.5% Ages 0-19	13.1% Ages 0-19	50.0% Ages 0-19		
		1.7% Ages 20+ CDRI 2015		2.2% Ages 20+ CDRI 2016	3.4% Ages 20+ CDRI 2017	4.2% Ages 20+ CDRI 2018	5.0% Ages 20+ CCAI		
Goal 12: Increase provider utilization of evidence-based treatment guidelines									
	There are no measures established for this goal. Refer to the Idaho Comprehensive Cancer Strategic Plan for objectives and strategic actions.								

## Quality of Life

Goal 13: Improve the physical and mental health of cancer survivors									
Indicator	Measure	Baseline	January 2017	October 2017	March 2019	June 2020	2020 Target	Progress towards target	Target Met
13.1	Proportion of cancer patients receiving survivorship care plans	TBA CoC Hospitals 2015					90.0% CoC		
13.2	Percentage of cancer survivors who report poor physical health 14+ of last 30 days (age adjusted to the year 2000 standard population)	29.0% BRFSS 2011-2012	20.5% BRFSS 2013-2014	16.5% BRFSS 2016	21.1% BRFSS 2017	22.7% BRFSS 2018	26.1% CCAI (10%)		✓
13.3	Percentage of cancer survivors who report poor mental health 14+ of last 30 days (age adjusted to the year 2000 standard population)	28.6% BRFSS 2011-2012	25.0% BRFSS 2013-2014	11.5% BRFSS 2016	16.5% BRFSS 2017	19.1% BRFSS 2018	25.7% CCAI (10%)		✓
13.4	Percentage of cancer survivors who are current smokers (age adjusted to the year 2000 standard population)	31.3% BRFSS 2011-2012	36.7% BRFSS 2013-2014	28.9% BRFSS 2016	19.9% BRFSS 2017	26.2% BRFSS 2018	28.1% CCAI (10%)		✓
13.5	Percentage of cancer survivors who report no physical activity outside of work (age adjusted to the year 2000 standard population)	25.9% BRFSS 2011-2012 (rev)	16.8% BRFSS 2013-2014	17.3% BRFSS 2016	22.3% BRFSS 2017	25.7% BRFSS 2018	23.3% CCAI (10%)		■
13.6	Percentage of cancer survivors who report consuming 5+ servings fruit and vegetables per day (age adjusted to the year 2000 standard population)	20.0% BRFSS 2011-2012	18.0% BRFSS 2013-2014		28.2% BRFSS 2017		22.0% CCAI (10%)		✓

Goal 14: Improve access and referrals to palliative care services for cancer patients									
Indicator	Measure	Baseline	January 2017				2020 Target	Progress towards target	Target Met
14.1	Access to palliative care in hospitals	B Grade CAPC 2015					A Grade CCAI		
14.2	Proportion of cancer patients whose physical pain caused by cancer or cancer treatment is under control	TBA BRFSS (n<50)							