


COLORECTAL CANCER in IDAHO

2023

Incidence	3,351 cases of malignant colorectal cancer were diagnosed among Idahoans during 2016-2020	Colorectal cancer rates are 40% higher in Idaho communities with higher rates of poverty. Colorectal cancer rates are 25% higher among Black Idahoans than among white Idahoans.
Mortality	1,318 Idahoans died of colorectal cancer during 2017-2021	Let's beat colorectal cancer! <ul style="list-style-type: none">• 45-75 is the recommend age to receive colorectal cancer screenings. Healthcare provider recommendation is the primary motivator for Idahoans to get screened.• Early detection saves lives! Colorectal cancer can be prevented or treated if found early.• Choose the best screening method – the one that gets done! Patients have several screening choices, including simple take-home options.
Prevalence	Over 3,700 Idahoans alive on July 1, 2020 had a colorectal cancer diagnosis within the prior 10 years	
Health Behavior	<ul style="list-style-type: none">• 1 in 3 eligible Idahoans are <u>not</u> up to date on their colorectal cancer screening (2020 BRFSS)• Idaho ranks 48th among U.S. states, the District of Columbia, and Puerto Rico for the proportion of adults aged 50-75 who are up to date on colorectal cancer screening (2020 BRFSS)	



Of every 20 Idahoans diagnosed with colorectal cancer, 7 have died from their cancer within 5 years of diagnosis

Rates of colorectal cancers have decreased **1.2%** per year since 2000.

In Idaho, males die from colorectal cancer at **1.3 times** the rate of females.

Even among tumors diagnosed at an early stage, **Hispanic Idahoans have ~10% lower 5-year survival post-diagnosis than non-Hispanic white Idahoans.**