RYAN CRENSHAW, M.D. Instructions for Colonoscopy with MoviPrep

Please read this packet, in its entirety, at least 2 weeks prior to your procedure. If you fail to follow the instructions and the procedure has to be cancelled, the cancellation fee will be charged.

If you need to cancel your procedure, please let us know at least <u>5 business days</u> prior to the procedure. If you fail to do so, you will be charged a \$250.00 cancellation fee.

You will receive a confirmation call from our office staff at least <u>5 days</u> prior to your scheduled procedure(s). All procedure(s) must be confirmed in order to remain on the schedule. If you do not receive a call from our office, please contact us at 703-444-4799.

If you take blood thinners such as Aspirin, Plavix, Xarelto or Coumadin, Dr. Crenshaw may recommend for you to hold these medications anywhere from 2-8 days prior to your procedure, depending on what agent you are taking.

Patients on Coumadin (Warfarin): If approved by the prescribing physician (i.e. cardiologist, neurologist or primary care provider), you will be asked to stop your Coumadin 5 days prior to your procedure(s). You will also be asked to obtain a PT/INR, PTT blood test the day prior to your procedure(s). If you did not receive an order for this blood test, please contact our office at (703) 444-4799. Dr. Crenshaw will instruct you regarding the date to restart Coumadin (Warfarin) on the day of your procedure(s).

Patients on Plavix (Clopidogrel): If approved by the prescribing physician (i.e. cardiologist, neurologist or primary care provider), you will be asked to stop the Plavix (clopidrogel) 7 days prior to your procedure(s). Dr. Crenshaw will instruct you regarding the date to restart Plavix (Clopidogrel) on the day of your procedure(s).

Patients taking Aspirin: Please make sure one of the two lines is checked off below. If not, please contact our office at (703) 444-4799.

 $_$ Please continue to take Aspirin, 81 mg or 325 mg, daily, including the day of the colon	oscopy.
Please stop taking Aspirin, 81 mg or 325 mg, 8 days prior to the colonoscopy,	

Any patient stopping Aspirin, Plavix, Coumadin or any other blood thinner should contact the prescribing doctor (primary care physician, or cardiologist) to confirm that it is acceptable to stop this medication(s) for the recommended period of time. You may take Tylenol if needed. Please do not take any other medication or products that can thin the blood, such as Ibuprofen, Motrin, Advil, Aleve, Vitamin E and/or Garlic pills 8 days prior to the procedure(s).

It is highly recommended that you take your medication for heart disease, high blood pressure and asthma every day, including the day of your procedure. If you take any medication(s) around the time that you are taking a dose of the laxative to prepare for your colonoscopy, please take the laxative first, then your medication. When taking medication(s) on the day of your procedure, the medication(s) must be taken at least 4 hours prior to your procedure time with water. At 4 hours prior to procedure time, you should stop consuming liquids, stop taking medications and not take anything orally until your procedure has been completed. All other medications should be brought to the hospital to be taken after your procedure.

If you are taking medications for diabetes, consult with the medical provider that is managing your diabetes to inform him/her that you are being asked to change your diet in preparation for colonoscopy. Please ask this provider how you should change your diabetes medication regimen to reduce the risk of your blood sugar becoming too low or too high during your preparation for the procedure.

You will need someone to drive you home from the hospital, or surgical center, after your procedure. You should not drive until the next day.

Preparation instructions for MoviPrep

(Sodium Sulfate, Sodium Chloride, Potassium Chloride, Sodium Ascorbate and Ascorbic Acid)

Caution: This preparation must be used with caution and may be contraindicated in patients with the following conditions: Kidney failure or compromise in kidney function, heart arrhythmias, history of seizure, impaired gag reflex, severe ulcerative colitis, esophageal regurgitation and/or gout. If you have any of these conditions, please call the office to confirm this preparation is appropriate for you.

Five days prior to the procedure: Do NOT eat food containing seeds, corn, nuts, black pepper, lettuce, raw vegetables, fruits with seeds or skin as they can be difficult to lavage from the colon. Please do not take fish oil, krill oil, lemon oil or any supplements or foods that contain a significant amount of oil such as potato chips.

The day prior to your procedure, you will be on a clear liquid diet (no solid food, except for Jello) for the entire day beginning with your breakfast meal. On the day before your colonoscopy, the more clear liquids you consume, the better for your preparation. In addition, adequate hydration will reduce the risk of developing headaches, lightheadedness and dizziness which can occur during the preparation.

It is also important to vary the liquids you are consuming. Do NOT restrict yourself to drinking water only. Please make sure you consume liquids with salt such as soup broth. And liquids with carbohydrates/sugars such as apple juice. By varying the liquids consumed, you reduce the risk of developing electrolyte abnormalities such as a low sodium. Please avoid any food or beverage product(s) which contain red or purple coloring.

MENU FOR CLEAR LIQUID DIET

<u>Breakfast</u>	<u>Lunch</u>	<u>Dinner</u>
White Cranberry Juice	Chicken broth	Chicken broth
Gelatin dessert	Apple Juice	White Grape Juice
Tea/coffee (no milk)	Sprite, 7up, Ginger Ale	Gelatin dessert
Gatorade or similar sports	Fruit-flavored ice	Sprite, 7up, Ginger Ale
drink	Tea/coffee (no milk)	Tea/coffee (no milk)

MOVIPREP MEDICATION
Instructions continue on the next page

FIRST DOSE: One day prior to your procedure you will need to mix the MoviPrep solution.

Empty 1 pouch A and 1 pouch B into the disposable container.

Add lukewarm drinking water to the top line of the container.

Cap bottle and mix to dissolve and refrigerate if desired.

You may add a tea bag to alter/enhance the taste if desired.

At 6:00 pm you will begin drinking the solution. The MoviPrep container is divided into 4 marks. Every 15 minutes, drink the solution down to the next mark which is equal to 8 ounces until the full liter is complete. Once you have finished the MoviPrep solution you may drink 16 ounces of a clear liquid of your choice from the clear liquid diet menu above.

SECOND DOSE: Day of the procedure: Please begin 7 hours prior to your procedure time.

You will need to mix the MoviPrep solution.

Empty 1 pouch A and 1 pouch B into the disposable container.

Add lukewarm drinking water to the top line of the container.

Cap bottle and mix to dissolve and refrigerate if desired.

You may add a tea bag to alter/enhance the taste if desired.

The MoviPrep container is divided by 4 marks. Every 15 minutes, drink the solution down to the next mark, until the full liter is completed.

Next drink 16 ounces of a clear liquid of your choice from the clear liquid diet menu listed above.

After completing the MoviPrep solution, you can consume up to 24 ounces of clear liquids over the next 60 minutes. You must be done with the clear liquids 4 hours prior to your scheduled procedure time. Please note that consuming any liquids or medications(s) too close to your procedure may result in delaying or cancelling your procedure. Do not consume any liquids or medication(s) by mouth for 4 hours prior to your procedure time.

On average, your bowel movements should be clear (clear = yellow or white liquid without solid or granular material) 3 hours after you finish drinking the solution. However, delays in the onset of bowel movements and/or becoming clear can occur several hours after completing the solution. If you are not clear at 2 hours after taking the second dose on the day of your procedure, please follow the instructions written below.

If you are unable to complete and/or tolerate the preparation for colonoscopy, please follow these instructions:

Instructions continue on the next page

Purchase the following (no prescription necessary) and begin this preparation 1/2 hour after the last glass of solution taken.

- One bottle of Magnesium Citrate
- One bottle of fleet enema (without mineral oil, or any other type of oil).

Drink one bottle of Magnesium Citrate. Wait 2 hours. If your bowel movements are not clear or you could not tolerate the Magnesium Citrate, then proceed with Fleet Enema as follows:

- 1. Apply one Fleet Enema (without mineral oil, or any other type of oil) per rectum and wait 30 minutes.
- 2. If your bowel movements are still not clear, fill the same enema bottle with warm 4an4e from the faucet. Then administer a warm 4an4e enema per rectum every 30 minutes until you have bowel movements which 4an4e4to f clear yellow or clear 4an4e liquid. Do NOT exceed more 4an 4 enemas.

If you still encounter significant difficulties with your preparation, please contact our office at (703) 444-4799. If you are forwarded to voicemail, follow directions to contact the doctor on call.

RYAN P. CRENSHAW, M.D.

21135 WHITFIELD PLACE, SUITE 102, STERLING, VA 20165 (703) 444-4799

OPERATIVE REQUEST/CONSENT

1.	procedure(s) along with surgical	assistants selected by perforation, and need for	Crenshaw (the "Practitioner") to perform him: colonoscopy, possible biopsy, risk of cor surgery has been explained. The Practities "Patient"):	drug allergy, over
	Please print your name:			
2.	been expected may present the do more than that which was sp named Practitioner and his surg judgment will be effective in the and radiology. I further authoric	mselves. I recognize, the cified in paragraph #2 gical assistants perform eir attempt to heal and, ze Anesthesiologist to a	urse of this operation some other condition at if such conditions are discovered it will above. I therefore authorize and request such surgical procedures which in their before diagnose. This includes, but is not limit administer whatever anesthesia they feeling personnel feel such is required.	be necessary to that the above- est professional red to, pathology
3.	I fully understand that this oper is guaranteed.	ation, like any operatio	n, is accompanied by some degree of risk	and that no cure
4.	4. The nature of my (or the patient's) condition, the nature of the procedure(s) listed under paragraph #1 above risks involved and whatever other choices are available to me (or the patient), if any, have been explained by the Practitioner. I have been given the opportunity to ask any questions that I may have regarding explanation and my questions have been answered satisfactorily.			explained to me
5.	I am aware of the "Cancellation not provided at least <u>5 business</u>		I that I will be held responsible for a \$250.0 eduled date for procedure.	00 fee if notice is
6.		•	ith both verbal and written instructions giv your procedure), you will be charged the	
7.	procedure(s), as well as obtaining	g the necessary referra	nsurance provider to check coverage fo als. It is also the patient's responsibility to vise, the patient will be held responsible fo	notify our office
Sig	nature of Patient	Date	Signature of Witness	Date.
	nature of Parent/Guardian nave personally explained, in non-	Date PHYSICIAN'S STATechnical terms, the pro	TEMENT oposed procedure to the patient, and/or F	Relative∕guardian,
	the major ris	sks or consequences of	this procedure, and any alternative.	

Page 5 of 5

PATIENT COPY: MoviPrep

RYAN P. CRENSHAW, M.D.

21135 WHITFIELD PLACE, SUITE 102, STERLING, VA 20165 (703) 444-4799

OPERATIVE REQUEST/CONSENT

Sigr	nature of Parent/Guardian	Date PHYSICIAN'S STA	TEMENT		
Sig	nature of Patient	Date	Signature of Witness	Date	
7.	procedure(s), as well as obtaining	ng the necessary referr	insurance provider to check coverage for als. It is also the patient's responsibility to vise, the patient will be held responsible for	notify our office	
6. If your procedure is cancelled due to non-compliance with both verbal and written instructions given not complying with the clear liquid diet the day prior to your procedure), you will be charged the ca					
5.	The nature of my (or the patient's) condition, the nature of the procedure(s) listed under paragraph #1 above, the risks involved and whatever other choices are available to me (or the patient), if any, have been explained to me by the Practitioner. I have been given the opportunity to ask any questions that I may have regarding that explanation and my questions have been answered satisfactorily. I am aware of the "Cancellation Policy" and understand that I will be held responsible for a \$250.00 fee if notice is not provided at least 5 business days in advance of scheduled date for procedure.				
:4.					
3.	I fully understand that this ope is guaranteed.	ration, like any operatio	on, is accompanied by some degree of risk	and that no cure	
2.	been expected may present the do more than that which was s named Practitioner and his sur judgment will be effective in the and radiology. I further author	emselves. I recognize, t pecified in paragraph # gical assistants perforn eir attempt to heal and ize Anesthesiologist to	urse of this operation some other condition hat if such conditions are discovered it will above. I therefore authorize and reques a such surgical procedures which in their for diagnose. This includes, but is not limical administer whatever anesthesia they fee ing personnel feel such is required.	II be necessary to it that the above- best professional ted to, pathology	
	Please print your name:		· · · · · · · · · · · · · · · · · · ·		
1.	procedure(s) along with surgica	al assistants selected by perforation, and need f	him: <u>colonoscopy, possible biopsy, risk of</u> or surgery has been explained. The Practit	drug allergy, over	
1	I hereby request consent to	and authorize Dr	Crenshaw (the "Practitioner") to perfor	m the following	

I have personally explained, in non-technical terms, the proposed procedure to the patient, and/or Relative/guardian, the major risks or consequences of this procedure, and any alternative.

Signature of	f Physician
--------------	-------------

Date

DOCTOR COPY: MoviPrep