## Physicians to Children Immunization Schedule

Visit		Immunization(s)					
Birth		Нер В	dose #1	given in the hospital or 1st office visit			
1 month		Нер В	dose #2	must have a minimum 28 days from dose #1			
2 month	Pentacel	Prevnar	Rotavirus***	must be a minimum of 6 weeks old			
4 month	Pentacel	Prevnar	Rotavirus***	minimum 6 weeks from 2 month immunizations			
6 month	Pentacel	Prevnar	Rotavirus***	minimum 6 weeks from 4 month immunizations			
9 month		Нер В	dose #3	must be a minimum of 6 months old			
12 months	MMR	Varivax	Нер А		must be 12 months old MMR can be given as early as 6 months, but will not count toward the series		
15 months	Pentacel	Prevnar			must be 15 months old		
18 months		Hep A **	dose #2	minimu	imum 6 months between Hep A doses		
2 year		Hep A **	dose #2	if no	not received at the 18 month visit		
4 year	Quadracel	Proquad (MMRV)			must be 4 years old		
11/12 year	TdaP	Menectra/ Menquadfi	HPV				
9-14 year	HPV #1	If started before age 15, only 2 doses are needed.					
	HPV #2	give 2nd dose 6 months after dose #1 or ideally not longer than 12 months					
15 year		HPV**	**3 doses needed if series initiated after 15th birthday. dose #2 is given 2 months after first doseHPV**dose #3 is given 4 months after second dose and 6 months from first dose				
16 year	Men B #1 (Bexsero)	Menactra/ Menquadfi #2	Menactra/Menquadifi - dose #2 is given 3-5 years from dose #1				
		Men B (Bexsero)	dose #2	minimum 30 days from dose #1			
Nursing Key		***Cannot start Rotavirus series if chil			Combination Vaccines		
No Color =	IM	over 14 weeks and 6 days old from the series can be given af			Pentacel =	DTaP, HIB & Polio	
	SubCut			ter child is 8	Prevnar =	Pneumococcal 13	
	Oral		months old.		Proquad =	MMR & Varicella	
					Quadracel=	DTaP & Polio	