

Surname	Centre Number	Candidate Number
Other Names		2



GCE A level

1095/01

CHEMISTRY – CH5

Section A	For Examiner’s use only		
	Question	Maximum Mark	Mark Awarded
	1.	10	
	2.	12	
	3.	18	
Section B	4.	20	
	5.	20	
	Total	80	

1095
010001

ADDITIONAL MATERIALS

In addition to this examination paper, you will need:

- a calculator;
- an 8 page answer book;
- a copy of the **Periodic Table** supplied by WJEC.
Refer to it for any **relative atomic masses** you require.

INSTRUCTIONS TO CANDIDATES

Use black ink or black ball-point pen.

Write your name, centre number and candidate number in the spaces at the top of this page.

Section A Answer **all** questions in the spaces provided.

Section B Answer **both** questions in **Section B** in a separate answer book which should then be placed inside this question-and-answer book.

Candidates are advised to allocate their time appropriately between **Section A (40 marks)** and **Section B (40 marks)**.

INFORMATION FOR CANDIDATES

The number of marks is given in brackets at the end of each question or part-question.

The maximum mark for this paper is 80.

Your answers must be relevant and must make full use of the information given to be awarded full marks for a question.


You are reminded that marking will take into account the Quality of Written Communication in all written answers.

SECTION A

Answer **all** questions in the spaces provided.

1. Ammonium salts are very important chemicals as they are used as a nitrogen source in fertilisers.


- (a) When cold aqueous sodium hydroxide is added to an ammonium salt, the following equilibrium exists.



Identify the **two** acid-base conjugate pairs in the equilibrium.

[2]

- (b) Ammonium chloride and sodium nitrite react together in aqueous solution to produce nitrogen gas. This can be represented by the ionic equation:



The rate equation for the reaction is given below.

$$\text{Rate} = k[\text{NH}_4^+][\text{NO}_2^-]$$

- (i) Complete the table of data for the above reaction. All experiments were carried out at the same temperature.

[3]

	$[\text{NH}_4^+(\text{aq})]/\text{mol dm}^{-3}$	$[\text{NO}_2^-(\text{aq})]/\text{mol dm}^{-3}$	Initial rate/ $\text{mol dm}^{-3} \text{s}^{-1}$
1	0.200	0.010	4.00×10^{-7}
2		0.010	2.00×10^{-7}
3	0.200		1.20×10^{-6}
4	0.100	0.020	

- (ii) Calculate the value of the rate constant, k , giving its units.

[2]

Value of k =

Units

- (iii) State how the value of k will alter, if at all, if the concentration of NH_4^+ ions is increased. [1]

.....

- (iv) State, giving a reason, how the value of k will alter, if at all, if the temperature is increased. [2]

.....

.....

.....

Total [10]

10

2. (a) Write an expression for the ionic product of water, K_w , giving its units, if any. [2]

$$K_w =$$

Units

- (b) (i) The value for K_w at 298 K is 1.0×10^{-14} . Explain why the pH of pure water at this temperature has a value of 7. [2]

.....

.....

.....

- (ii) Calculate the pH of the final solution if 10 cm^3 of 0.10 mol dm^{-3} hydrochloric acid is added to 990 cm^3 of pure water. [2]

pH =

- (c) Calculate the pH of a solution which is $0.010 \text{ mol dm}^{-3}$ with respect to ethanoic acid and $0.020 \text{ mol dm}^{-3}$ with respect to sodium ethanoate at 298 K. [3]

[K_a for ethanoic acid = $1.78 \times 10^{-5} \text{ mol dm}^{-3}$ at 298 K]

pH =

- (d) If 10 cm^3 of 0.10 mol dm^{-3} hydrochloric acid is added to 990 cm^3 of the solution described in (c) the change in pH is only 0.06. Explain why this change in pH is much smaller than that in (b)(ii). [3]

.....

.....

.....

.....

.....

.....

Total [12]


12

3. Read the passage below and then answer the questions in the spaces provided.

Hydrogen Peroxide

If a non-scientist knows only one chemical formula it is most likely to be H_2O for water but how much do you know about another hydrogen oxide, hydrogen peroxide? A molecule of hydrogen peroxide has the molecular formula H_2O_2 .


- 5 Most chemistry students first meet hydrogen peroxide as a colourless solution that is used to prepare oxygen. Bottles of hydrogen peroxide from a pharmacist are often labelled '20 volume'. This means that one volume of solution decomposes to give 20 volumes of oxygen gas. The equation for the decomposition is:



- 10 This reaction is very slow at room temperature. However the addition of a suitable catalyst increases the rate of decomposition phenomenally. Manganese(IV) oxide, potatoes and blood are all effective. Potatoes and blood both contain the enzyme catalase and one catalase molecule decomposes 50 000 molecules of H_2O_2 per second!


Is hydrogen peroxide an oxidising agent or a reducing agent?

- 15 Both in the laboratory and at home hydrogen peroxide is most commonly used as an oxidising agent (so the hydrogen peroxide itself is reduced). The half-equation is:




Since some colouring matter is bleached by oxidation and the product of hydrogen peroxide's reduction is water, it is used as a safe bleaching agent particularly in hair treatment. A peroxide blonde is someone with almost white hair, usually as a result of treatment with hydrogen peroxide.

- 20 However, if hydrogen peroxide reacts with a more powerful oxidising agent such as potassium manganate(VII), the hydrogen peroxide will act as a reducing agent and will itself be oxidised. The half-equation is:



- 25 Therefore hydrogen peroxide can act as both oxidising agent and reducing agent. In fact, it can react with itself so that alternate molecules are oxidised and reduced. The overall equation is obtained by adding the half-equations for the reduction and oxidation, giving



which is the standard decomposition equation!

- End of passage -

- (a) Using outer electrons only, draw a dot and cross diagram to show the bonding in a hydrogen peroxide molecule (*line 3*). [1]

- (b) Use the equation for the decomposition of hydrogen peroxide (*line 8*) to calculate the concentration, in mol dm^{-3} , of aqueous hydrogen peroxide solution in a bottle of '20 volume hydrogen peroxide' at 25°C . [2]

[1 mol of oxygen occupies 24 dm^3 at 25°C]

Concentration = mol dm^{-3}

- (c) Manganese(IV) oxide (*line 10*) and potassium manganate(VII) (*lines 20-21*) are typical transition metal compounds.

- (i) Give **two** reasons why transition metal compounds can act as catalysts. [2]

.....

.....

.....

- (ii) Explain why transition metal complex ions appear coloured. [4]
QWC [1]

.....

.....

.....

.....

.....

.....

.....

.....

(d) In an acidic solution, hydrogen peroxide is oxidised to oxygen by potassium manganate (VII) (*lines 20-23*).

(i) Write the half-equation for the reduction of MnO_4^- to Mn^{2+} ions in acidic solution.

[1]

(ii) Use your answer to (i) and the half-equation given in *line 23* to deduce the overall equation for this reaction.

[2]

(iii) 20.0 cm^3 of an acidified solution of hydrogen peroxide required 14.80 cm^3 of a 0.020 mol dm^{-3} solution of potassium manganate(VII) for complete reaction. Calculate the concentration, in mol dm^{-3} , of the hydrogen peroxide solution.

[3]

Concentration = mol dm^{-3}

(e) Explain, using oxidation states, why the decomposition of hydrogen peroxide (*line 27*) can be classified as a redox reaction.

[2]

Total [18]


18

Total Section A [40]

SECTION B

Answer **both** questions in the separate answer book provided.

4. (a) The diagram shows some of the reactions of lead compounds.



- (i) State the role of lead(IV) oxide in the reaction with concentrated hydrochloric acid. [1]
- (ii) Name white solid **A** and gas **B**. [2]
- (iii) Give the formula of the lead-containing species present in colourless solution **D**. [1]
- (iv) Give the colour of precipitate **E**. [1]
- (v) Write the equation for the formation of lead(II) nitrate from lead(II) oxide. [1]

(b) Carbon is the first element in Group 4. Two of its allotropes are diamond and graphite. A compound that forms structures corresponding to diamond and graphite is boron nitride.

(i) Describe the structure of graphite and explain why **hexagonal** boron nitride can adopt the same structure yet have different electrical conductivity properties. [4]
QWC [1]

(ii) State **one** use for the **cubic** boron nitride structure. [1]

(c) Another element in Group 4 is tin. At low temperatures tin exists as its grey form. At higher temperatures the white form is stable. The change can be represented by the equation:



The standard entropy values are $44.8 \text{ J K}^{-1} \text{ mol}^{-1}$ for grey tin and $51.5 \text{ J K}^{-1} \text{ mol}^{-1}$ for white tin.

(i) Calculate the minimum temperature needed to cause grey tin to change to white tin. [3]

(ii) During Napoleon's disastrous campaign in Russia from June to December in 1812 the tin buttons on his infantry's uniforms disintegrated. Suggest a reason why this might have happened. [1]

(d) An important technological development in recent years has been the hydrogen fuel cell. This uses electrochemical methods to get energy from hydrogen.

(i) Write the half-equations for the processes occurring at the electrodes and an equation for the overall reaction. [3]

(ii) Give **one** disadvantage of using hydrogen fuel cells to power vehicles. [1]

Total [20]

5. (a) Chlorine reacts with aqueous sodium hydroxide in one of two ways, depending on the temperature used.

(i) Write the equation for the reaction of chlorine with

I cold aqueous sodium hydroxide, [1]

II hot aqueous sodium hydroxide. [1]

(ii) Classify this type of redox reaction. [1]

- (b) Chlorine reacts with many elements to form chlorides. Explain why phosphorus forms two chlorides, PCl_3 and PCl_5 , but nitrogen only forms NCl_3 . [2]

- (c) Most ionic chlorides, e.g. sodium chloride, are soluble in water. However some, e.g. silver chloride, are insoluble.

The enthalpy change of solution of an ionic compound and its solubility depend on the balance between two enthalpy changes. Name these enthalpy changes and state if they are endothermic or exothermic. Explain how the enthalpy change of solution of a compound and its solubility depend on the balance between them. [4]

QWC [1]


- (d) Some standard electrode potentials, E^\ominus , are given below.

System	E^\ominus / V
$\frac{1}{2} \text{I}_2(\text{s}) + \text{e}^- \rightleftharpoons \text{I}^-(\text{aq})$	+0.54
$\text{Fe}^{3+}(\text{aq}) + \text{e}^- \rightleftharpoons \text{Fe}^{2+}(\text{aq})$	+0.77
$\frac{1}{2} \text{Br}_2(\text{l}) + \text{e}^- \rightleftharpoons \text{Br}^-(\text{aq})$	+1.09
$\frac{1}{2} \text{Cl}_2(\text{g}) + \text{e}^- \rightleftharpoons \text{Cl}^-(\text{aq})$	+1.36
$\text{Ce}^{4+}(\text{aq}) + \text{e}^- \rightleftharpoons \text{Ce}^{3+}(\text{aq})$	+1.45

- (i) Using the information from the table, state which of the **halides** will reduce Fe^{3+} to Fe^{2+} . Give a reason for your answer. [2]
- (ii) Write the cell diagram of the cell formed by combining the $\text{Fe}^{3+}(\text{aq})$, $\text{Fe}^{2+}(\text{aq})$ and $\text{Ce}^{4+}(\text{aq})$, $\text{Ce}^{3+}(\text{aq})$ half cells and calculate the standard e.m.f. of this cell. [2]

QUESTION 5 CONTINUES ON PAGE 12

- (e) A flask containing an initial mixture of 0.100 mol of ethanoic acid and 0.083 mol of methanol was kept at 25 °C until the following equilibrium had been established.



The ethanoic acid present at equilibrium required 32.0 cm³ of a 1.25 mol dm⁻³ solution of sodium hydroxide for complete reaction.

- (i) Write an expression for the equilibrium constant, K_c , giving the units, if any. [2]
- (ii) Calculate the number of moles of ethanoic acid present at equilibrium. [1]
- (iii) Calculate the value of the equilibrium constant, K_c , for this reaction. [2]
- (iv) State, giving a reason, what happens to the value of the equilibrium constant, K_c , if the temperature is increased. [1]

Total [20]

Total Section B [40]

END OF PAPER



GCE A level

1095/01-A

CHEMISTRY – CH5

Periodic Table

P.M. TUESDAY, 17 June 2014

