

Articles



Canine Demodicosis

Spring 2005

Signalment and History

“Buffy”, a 10 year old spayed female Shih Tzu, presented for a 6 month history of mildly pruritic patchy alopecia and scaling of the face, legs and truncal regions which had failed to respond to several courses of antibiotics or steroid injections. She had also become lethargic the last few months.

Differential Diagnoses

The patchy alopecia and scaling were consistent with a folliculitis (bacterial infection, demodicosis, dermatophytosis), and/or malassezia dermatitis possibly due to an underlying endocrine disease (hypothyroidism, iatrogenic or endogenous hyperadrenocorticism) or hypersensitivity (atopy or food allergy). Less common causes of alopecia and scaling included neoplasia (cutaneous T cell lymphoma), and sebaceous adenitis.



Laboratory Findings

A dry skin scraping for cytology revealed occasional neutrophils with scattered intracellular and extracellular cocci bacteria. A dermatophyte culture was negative. Deep skin scrapings revealed demodex mites in all life stages (adults, larvae, nymphs and eggs). Full bloodwork was performed. The complete blood count revealed mild normocytic, normochromic anemia. The chemistry panel revealed mildly elevated ALP consistent with prior

steroid therapy and a mild hyperglobulinemia. A thyroid profile revealed a low total T4, low free T4 by equilibrium dialysis and a high TSH level consistent with the diagnosis of hypothyroidism.

Treatment

Buffy was started on cephalexin (22 mg/kg PO BID for 3 weeks) to treat the bacterial folliculitis. Thyroid supplementation was initiated with Soloxine at (0.1 mg/10 lbs PO BID). The demodicosis was treated with 1% ivermectin at a gradually increasing dosage regimen beginning at 0.1 mg/kg PO once daily over 2 weeks reaching a final treatment dosage of 0.5 mg/kg PO once daily.



Follow Up

Found in hair follicles, and an unnamed short-bodied Demodex mite which lives in the stratum corneum are rarer causes of skin disease in dogs.

In dogs, demodex infections can present as localized (more common in puppies 3-6 months old) or generalized (juvenile-onset or adult-onset) disease. Juvenile-onset generalized demodicosis occurs in young dogs, between 3-18 months of age, most commonly in medium to large purebred dogs. Adult-onset generalized demodicosis occurs in dogs greater than 18 months of age, with the highest incidence in middle-aged to older dogs that are immunocompromised due to an underlying condition such as hypothyroidism, iatrogenic or acquired hyperadrenocorticism, diabetes mellitus, immunosuppressive drug therapy or neoplasia. Demodicosis lesions begin as patchy alopecia with variable erythema, scaling, and papules. Pruritus is variable. With chronicity, the affected skin may become hyperpigmented, lichenified, pustular, eroded and/or ulcerated from a secondary superficial or deep pyoderma. Lesions can occur anywhere on the body, including the feet, and severe pododermatitis is not uncommon. Peripheral lymphadenopathy can be observed. Differential diagnoses for canine demodicosis include superficial or deep pyoderma, dermatophytosis, hypersensitivity (flea allergy, food allergy, atopy), and pemphigus foliaceus.

Diagnosis of demodicosis

Seems to have a genetic basis in dogs, so affected animals should not be bred. Localized demodex lesions may resolve spontaneously. If necessary, localized lesions in dogs can be treated with Rotenone (Canex) or 0.025% amitraz solution daily. Traditional miticidal therapy for generalized demodicosis in dogs entails clipping medium- to long-coated breeds, bathing in 2.5-3% benzoyl peroxide shampoo, and then following up with a total body application of 0.025%-0.05% amitraz solution weekly (Cure rate 50-86%).



Additional treatment options for generalized canine demodicosis include ivermectin (except in Collies, Shelties, herding breeds, and white German Shepherds, breeds which have an increased risk of having a genetic defect enabling penetration of the blood-brain barrier by ivermectin). 1% ivermectin is given orally beginning at a dose of 0.1 mg/kg/day and increased by 0.1 mg/kg every 3 days assuming toxicity does not occur. The therapeutic dosage range is from 0.3-0.6 mg/kg/day. A cure rate of 85-90% can be achieved at 0.6 mg/kg/day. Expect to treat cases for 3-6 months. Some sensitive animals will experience signs of toxicity including mydriasis, blindness, ataxia, dullness or seizures, and if toxic signs occur the drug must be immediately discontinued. Prior to instituting ivermectin therapy in a potentially sensitive dog, testing for the MDR-1 genetic defect is now available at Washington State University (www.vetmed.wsu.edu/depts-VCPL).

An alternative treatment option, which is safe for the ivermectin-sensitive breeds but more expensive, is milbemycin oxime (Interceptor) 1-2 mg/kg orally once daily. All the above mentioned treatments should be continued until 2 separate skin scrapings, 30 days apart, are negative (no live or dead mites). Relapses may occur in up to 15% of cases and require pulse or daily long-term treatment for control. Lime sulfur dips, selamectin, and once weekly ivermectin injections are ineffective for treatment of generalized canine demodicosis. The prognosis for localized demodicosis is good, while for generalized disease the prognosis varies from good to guarded depending on the underlying cause.

Suggested readings:

Scott D, Miller M, Griffin C.

Small Animal Dermatology, 6th edition. W.B. Saunders Company, Philadelphia, PA 2001. pp 457-476.

Hnilica K, Medleau L.

Small Animal Dermatology: A Color Atlas and Therapeutic Guide. W.B. Saunders Company. Philadelphia, PA 2001. pp 65-70.