RYAN CRENSHAW, M.D. Instructions for Colonoscopy with SuPrep

Please read this packet, in its entirety, at least 2 weeks prior to your procedure. If you fail to follow the instructions and the procedure has to be cancelled, the cancellation fee will be charged.

If you need to cancel your procedure, please let us know at least <u>5 business days</u> prior to the procedure. If you fail to do so, you will be charged a \$250.00 cancellation fee.

You will receive a confirmation call from our office staff at least <u>5 days</u> prior to your scheduled procedure(s). All procedure(s) must be confirmed in order to remain on the schedule. If you do not receive a call from our office, please contact us at 703-444-4799.

If you take blood thinners such as Aspirin, Plavix, Xarelto or Coumadin, Dr. Crenshaw may recommend for you to hold these medications anywhere from 2-8 days prior to your procedure, depending on what agent you are taking.

Patients on Coumadin (Warfarin): If approved by the prescribing physician (i.e. cardiologist, neurologist or primary care provider), you will be asked to stop your Coumadin 5 days prior to your procedure(s). You will also be asked to obtain a PT/INR, PTT blood test the day prior to your procedure(s). If you did not receive an order for this blood test, please contact our office at (703) 444-4799. Dr. Crenshaw will instruct you regarding the date to restart Coumadin (Warfarin) on the day of your procedure(s).

Patients on Plavix (Clopidogrel): If approved by the prescribing physician (i.e. cardiologist, neurologist or primary care provider), you will be asked to stop the Plavix (clopidrogel) 7 days prior to your procedure(s). Dr. Crenshaw will instruct you regarding the date to restart Plavix (Clopidogrel) on the day of your procedure(s).

Patients taking Aspirin: Please make sure one of the two lines is checked off below. If not, please contact our office at (703) 444-4799.

 Please continue to take Aspirin,	, 81 mg or 325 mg,	daily, including the da	y of the colonoscopy.
Please stop taking Aspirin, 81 r	ng or 325 mg, 8 da	rys prior to the colonos	copy,

Any patient stopping Aspirin, Plavix, Coumadin or any other blood thinner should contact the prescribing doctor (primary care physician, or cardiologist) to confirm that it is acceptable to stop this medication(s) for the recommended period of time. You may take Tylenol if needed. Please do not take any other medication or products that can thin the blood, such as Ibuprofen, Motrin, Advil, Aleve, Vitamin E and/or Garlic pills 8 days prior to the procedure(s).

It is highly recommended that you take your medication for heart disease, high blood pressure and asthma every day, including the day of your procedure. If you take any medication(s) around the time that you are taking a dose of the laxative to prepare for your colonoscopy, please take the laxative first, then your medication. When taking medication(s) on the day of your procedure, the medication(s) must be taken at least 4 hours prior to your procedure time with water. At 4 hours prior to procedure time, you should stop consuming liquids, stop taking medications and not take anything orally until your procedure has been completed. All other medications should be brought to the hospital to be taken after your procedure.

If you are taking medications for diabetes, consult with the medical provider that is managing your diabetes to inform him/her that you are being asked to change your diet in preparation for colonoscopy. Please ask this provider how you should change your diabetes medication regimen to reduce the risk of your blood sugar becoming too low or too high during your preparation for the procedure.

You will need someone to drive you home from the hospital, or surgical center, after your procedure. You should not drive until the next day.

Preparation Instructions for SuPrep (Sodium sulfate, Potassium sulfate, and Magnesium sulfate)

Caution: This preparation must be used with caution and may be contraindicated in patients with the following conditions: Kidney failure or compromise in kidney function, heart arrhythmias, history of seizures, impaired gag reflex, severe ulcerative colitis, esophageal regurgitation and/or gout. If you have any of these conditions, please call the office to confirm this preparation is appropriate for you.

Five days prior to the procedure: Do NOT eat food containing seeds, corn, nuts, black pepper, lettuce, raw vegetables, fruits with seeds or skin as they can be difficult to lavage from the colon. Please do not take fish oil, krill oil, lemon oil or any supplements or foods that contain a significant amount of oil such as potato chips.

The day prior to your procedure, you will be on a clear liquid diet (no solid food, except for Jello) for the entire day beginning with your breakfast meal. On the day before your colonoscopy, the more clear liquids you consume, the better for your preparation. In addition, adequate hydration will reduce the risk of developing headaches, lightheadedness and dizziness which can occur during the preparation.

It is also important to vary the liquids you are consuming. Do NOT restrict yourself to drinking water only. Please make sure you consume liquids with salt such as soup broth. And liquids with carbohydrates/sugars such as apple juice. By varying the liquids consumed, you reduce the risk of developing electrolyte abnormalities such as a low sodium. Please avoid any food or beverage product(s) which contain red or purple coloring.

MENU FOR CLEAR LIQUID DIET

Breakfast Lunch Dinner White Cranberry Juice Chicken broth Chicken broth Gelatin dessert Apple Juice White Grape Juice Sprite, 7up, Ginger Ale Gelatin dessert Tea/coffee (no milk) Gatorade or similar sports Fruit-flavored ice Sprite, 7up, Ginger Ale Tea/coffee (no milk) drink Tea/coffee (no milk)

SUPREP MEDICATION

DO NOT FOLLOW THE INSTRUCTIONS WRITTEN ON THE SUPREP BOX THAT CONTAIN THE BOTTLES OF MEDICINE. FOLLOW THE INSTRUCTIONS WRITTEN BELOW

Both 6 ounce bottles are required for a complete preparation.

Instructions continue on the next page

Page 2 of 4

FIRST DOSE: One day prior to your procedure, at 6:00 pm, complete steps 1 through 2

- Step 1 Pour one 6 ounce bottle of SuPrep liquid into the mixing container and add cool drinking water to the 16-ounce line on the container and mix. Drink all of the liquid in the container within one hour. If you need to take any medications at around this time, please take the dose of Suprep first, then take your medication.
- Step 2 You must drink 2 more 16-ounce containers of water over the next hour. You may then continue to consume clear liquids listed above.

SECOND DOSE: Day of the procedure: Please begin 6 hours prior to your procedure time.

Step 3 — Repeat steps 1 and 2 as written above. You must finish drinking the final glass of water at least 4 hours before your procedure time. If you need to take any medications at around this time, please take the dose of Suprep first, then take your medication. If you finish the Suprep and water recommended ahead of schedule, you may consume more clear liquids listed above until 4 hours prior to your procedure time. Please do not consume any liquids or medications after the time that corresponds to 4 hours before your procedure time.

On average, your bowel movements should be clear (clear = yellow or white liquid without solid or granular material) 3 hours after you finish drinking the solution. However, delays in the onset of bowel movements and/or becoming clear can occur several hours after completing the solution. If you are not clear at 2 hours after taking the second dose on the day of your procedure, please follow the instructions written below.

If you are unable to complete and/or tolerate the preparation for colonoscopy, please follow these instructions:

Purchase the following (no prescription necessary) and begin this preparation 1/2 hour after the last glass of solution taken.

- One bottle of Magnesium Citrate
- One bottle of fleet enema

Drink one bottle of Magnesium Citrate. Wait 2 hours. If your bowel movements are not clear or you could not tolerate the Magnesium Citrate, then proceed with Fleet Enema as follows:

- 1. Apply one Fleet Enema per rectum and wait 30 minutes.
- 2. If your bowel movements are still not clear, fill the same enema bottle with warm water from the faucet. Then administer a warm water enema every 30 minutes until you have bowel movements which consist of clear yellow or clear white liquid. Do NOT exceed more than 4 enemas.

If you still encounter significant difficulties with your preparation, please contact our office at (703) 444-4799. If you are forwarded to voicemail, follow directions to contact the doctor on call.

RYAN P. CRENSHAW, M.D.21135 WHITFIELD PLACE, SUITE 102, STERLING, VA 20165 (703) 444-4799

OPERATIVE REQUEST/CONSENT

1.	I hereby request, consent to, and authorize Dr. Crenshaw (the "Practitioner") to perform the following procedure(s) along with surgical assistants selected by him: colonoscopy, possible biopsy, risk of drug allergy, over sedation, aspiration, bleeding, perforation, and need for surgery has been explained. The Practitioner has advised me there is a small possibility of missing lesions on (the "Patient'):						
	Please print your name:						
2.	It has been clearly explained to me that during the course of this operation some other conditions that have not been expected may present themselves. I recognize, that if such conditions are discovered it will be necessary to do more than that which was specified in paragraph #1 above. I therefore authorize and request that the above-named Practitioner and his surgical assistants perform such surgical procedures which in their best professional judgment will be effective in their attempt to heal and/or diagnose. This includes, but is not limited to, pathology and radiology. I further authorize Anesthesiologist to administer whatever anesthesia they feel is indicated and authorize the use of blood transfusion(s) when attending personnel feel such is required.						
3.	I fully understand that this guaranteed.	operation, like any	operation, is ac	companied by some	e degree of risk and	that no cure is	
: 4.	The nature of my (or the pat involved and whatever other Practitioner. I have been give questions have been answere	r choices are availa en the opportunity t	ible to me (or t	he patient), if any,	have been explaine	ed to me by the	
5.	I am aware of the "Cancellat provided at least 5 business of	1 -		-	ble for a \$250.00 fe	e if notice is not	
6.	If your procedure is cancelle complying with the clear liqu	_			-	_	
7.	It is the patient's responsibility well as obtaining the necessary insurance provider changes, o	ry referrals. It is al	so the patient's	responsibility to no	otify our office imm	ediately if your	
Sig	nature of Patient	Date	<u></u>	Signature of W	itness	Date	
Sign	nature of Parent/Guardian	Date					
	I have personally explained, ir maj			procedure to the pa	· ·	e/guardian, the	
		Signature of	Physician	Date	NAMES NA		
		Page 4	of 4		PATIENT COPY:	SuPrep	

RYAN P. CRENSHAW, M.D.21135 WHITFIELD PLACE, SUITE 102, STERLING, VA 20165 (703) 444-4799

OPERATIVE REQUEST/CONSENT

		(the "Practitioner") to perform the followssible biopsy, risk of drug allergy,				
	d need for surgery has been	explained. The Practitioner has adv				
Please print your name:						
expected may present the than that which was speci- and his surgical assistants in their attempt to heal an Anesthesiologist to admir	clearly explained to me that during the course of this operation some other conditions that have not been by present themselves. I recognize, that if such conditions are discovered it will be necessary to do more that the specified in paragraph #1 above. I therefore authorize and request that the above-named Practitioner call assistants perform such surgical procedures which in their best professional judgment will be effective not to heal and/or diagnose. This includes, but is not limited to, pathology and radiology. I further authorize to administer whatever anesthesia they feel is indicated and authorize the use of blood transfusion(s) and personnel feel such is required.					
3. I fully understand that th guaranteed.	is operation, like any operation	n, is accompanied by some degree o	f risk and that no cure is			
involved and whatever of	her choices are available to make the opportunity to ask any	of the procedure(s) listed under para ne (or the patient), if any, have beer y questions that I may have regarding	explained to me by the			
	llation Policy" and understand ss days in advance of schedule	that I will be held responsible for a \$3 d date for procedure.	250.00 fee if notice is not			
	_	ith both verbal and written instruction procedure), you will be charged the				
well as obtaining the nece	ssary referrals. It is also the p	e provider to check coverage for the ratient's responsibility to notify our or held responsible for any charges for the	ffice immediately if your			
Signature of Patient	Date	Signature of Witness	Date			
ignature of Parent/Guardian	Date					
	_	ATEMENT coposed procedure to the patient, and/other this procedure, and any alternative.	or Relative/guardian, the			
	Signature of Physicia	n Date				
		DOCTOR CO	OPY: SuPrep			