

How to Get Rid of The Ill Effects of Radiation Poisoning



- Pour one pound of Epsom salts ... OR ... sea salt into a warm bath
- Add one pound of baking soda into a warm bath.
- Soak in the water for twenty minutes.
- Immerse the part affected if possible.
- Do this twice each week for one month.
- If you have any surgical scars or radiation burns, spread on YOUTH Moisture Activating Serum or Enfuselle Hand & Body Lotion. If the skin is NOT open, you can use Youth Radiance C+E



Youth Moisture
Activating Serum



Shaklee Hand and
Body Lotion



Radiance

Radiation Poisoning or Radiation Sickness:

Radiation exposure can also increase the probability of contracting some other diseases, mainly cancer, tumors, and genetic damage.

Mild radiation sickness

- Nausea and vomiting within 24 to 48 hours
- Headache
- Fatigue
- Weakness

Moderate radiation sickness

- Nausea and vomiting within 12 to 24 hours
- Fever
- Hair loss
- Infections
- Vomiting blood
- Bloody stool
- Poor wound healing
- Any signs and symptoms associated with a lower absorbed dose

Moderate radiation sickness can be fatal to those most sensitive to radiation exposure.

Severe radiation sickness

- Nausea and vomiting less than one hour after exposure to radiation
- Diarrhea
- High fever
- Any signs and symptoms associated with a lower absorbed dose
- Low blood pressure (hypotension)
- Severe radiation sickness is fatal about half the time.