Stone Curlew

| Burhinus oedicnemus | Category A |
|---------------------|------------|
| Very rare vagrant. | 2 records, |
| | 2 birds |

World/British distribution (Snow & Perrins 1998, Holling et al. 2019):

It has a scattered distribution mainly across southern Europe and North Africa, extending eastwards to central Asia. It is mainly migratory in northern and eastern Europe, but often varying from partially migratory to resident in milder areas. Substantial numbers of European birds cross the Mediterranean to winter in Africa. It has undergone a widespread and often severe contraction of its range, due mainly to loss of suitable breeding habitat through intensification of arable farming and conversion of dry grassland to cultivation. In Britain it is restricted mainly to East Anglia and Salisbury Plain, where there around 350 breeding pairs.

Kent status (KOS 2022):

It is now a rare passage migrant in Kent.

Latham (1785) wrote of the Stone Curlew as "not uncommon" in some parts of Kent, including the Downs but by Ticehurst's time (1909), the numbers were obviously much lower and breeding restricted to Dungeness, where a small population persisted until around 1970 (Taylor *et al.* 1981).

Knight and Tolputt (1871) included the "Great Plover" (as it was then known) in their "List of birds observed in Folkestone and its immediate neighbourhood". Whilst this range extends further inland that the current Folkestone and Hythe area, so is not conclusive evidence of occurrence here, it does suggest that the species was present locally, presumably on the Downs or possibly on the coastal shingle, in the days when it was more regular in the county.

In 1952 Roger Norman was informed by a Warrant Officer that one had been seen on the Hythe Ranges (in the vicinity of serial 18) in April of that year (Norman 2006).

There has only been one modern area record, which involved one found by Dale Gibson at Copt Point on the 6th September 1990, with presumably the same individual being relocated there on the 2nd and 3rd October. On one occasion it was seen roosting with gulls on the beach.

The modern record is shown by year in figure 1.




Figure 1: Stone Curlew records at Folkestone and Hythe since 1985




Figure 2: Stone Curlew records at Folkestone and Hythe by week

Figure 3 shows the location of records by tetrad.




Figure 3: Distribution of all Stone Curlew records at Folkestone and Hythe by tetrad

The full list of records is as follows:

1952 Hythe Ranges, one, in April (per R. K. Norman)
1990 Copt Point, one, 6th September, presumed same on 2nd to 3rd October (D. A. Gibson)

References

Holling, M. & the Rare Breeding Birds Panel 2019. Rare breeding birds in the UK in 2017. British Birds 112: 706-758.

Kent Ornithological Society (KOS) 1953-2022. Kent Bird Reports. Kent Ornithological Society.

Knight, V. & Tolputt, F. 1871. List of birds observed in Folkestone and its immediate neighbourhood. *Fourth Annual Report of the Folkestone Natural History Society*. Available at: <u>www.biodiversitylibrary.org/item/46103</u>

Latham, J. 1785. A General Synopsis of Birds. Benj. White, London.

Norman, R. K. 2006. Some observations of birds on Hythe Ranges since 1950. <u>https://folkestonebirds.com/where-to-watch</u> (see "Further Information" section).

Snow, D. & Perrins, C.M. 1998. The Birds of the Western Palearctic. Oxford University Press.

Taylor, D., Davenport, D. & Flegg, J. 1981. *Birds of Kent*. Kent Ornithological Society.

Ticehurst, N. F. 1909. *A History of the Birds of Kent*. Witherby, London.

Acknowledgements

The tetrad map images were produced from the Ordnance Survey <u>Get-a-map service</u> and are reproduced with kind permission of <u>Ordnance Survey</u>.

I am indebted to Andrew Henderson and Tim Hodge for providing access to the Kent Ornithological Society archives.